

RESONANCES AND SPECTRAL SHIFT FUNCTION FOR THE SEMI-CLASSICAL DIRAC OPERATOR

ABDALLAH KHOCHMAN

ABSTRACT. We consider the self-adjoint operator $H = H_0 + V$, where H_0 is the free semi-classical Dirac operator on $\mathbb{R}^3$. We suppose that the smooth matrix-valued potential $V = O(\langle x \rangle^{-\delta})$, $\delta > 0$, has an analytic continuation in a complex sector outside a compact. We define the resonances as the eigenvalues of the non-selfadjoint operator obtained from the Dirac operator H by a complex distortions of $\mathbb{R}^3$. We establish an upper bound $O(h^{-3})$ for the number of resonances in any compact domain. For $\delta > 3$, a representation of the derivative of the spectral shift function $\xi(\lambda, h)$ related to the semi-classical resonances of H and a local trace formula are obtained. In particular, if V is an electro-magnetic potential, we deduce a Weyl-type asymptotic of the spectral shift function. As a by-product, we obtain an upper bound $O(h^{-2})$ for the number of resonances close to non-critical energy levels in domains of width h and a Breit-Wigner approximation formula for the derivative of the spectral shift function.

Keywords: Semi-classical Dirac operator - Resonances - Trace formula - Spectral shift function - Weyl-type asymptotics - Breit-Wigner approximation.

Mathematics classification: 35B34 - 35P05 - 34L40 - 81Q20 - 81Q10.

1. INTRODUCTION

The resonance theory for the Schrödinger equation has been developed following several approaches. Among them we can mention the analytic dilation (see [1]) or the analytic distortion (see [21]) and, in the semi-classical regime, that related to the work of Helffer-Sjöstrand [20]. In [18] Helffer-Martinez showed that the different definitions give the same resonances when one can simultaneously apply them to an operator. For the three dimensional Dirac operator, Seba [41] defined the resonances as complex eigenvalues of the operator obtain by a complex dilation. Applying the approach of Helffer-Sjöstrand [20], Parisse [29] has studied the Dirac resonances in the semi-classical regime, with some scaling functions. The last two works deal with analytic perturbations near the real axis.

The concept of the spectral shift function has been introduced by Lifshits [25] in connection with the problems in quantum statistics and solid physics. Thereafter, a mathematical theory of the spectral shift function has been constructed by Krein [24]. Moreover, in [3] Birman-Krein found a connection between the scattering theory and the theory of the spectral shift function. A detailed presentation of the theory of the spectral shift function can be found in [44]. For a survey concerning the spectral shift function (SSF) for Schrödinger and Dirac operator or the asymptotic expansion of this function, we refer to Robert [35] and to the references given there.

A representation of the derivative of the scattering phase in terms of the resonances has been established for the Schrödinger operators. Such representations have been successively obtained by Melrose [26] for obstacle problems in the high energy case, by Petkov-Zworski [31], [32] for "black box" scattering with compact perturbations in the classical and the semi-classical cases and by Bruneau-Petkov [8] for long-range perturbations in the semi-classical "black box" framework. The results in [8] have been generalized by Dimassi-Petkov [13] for non-semi-bounded Schrödinger type operators. As a by-product, they prove a Weyl type asymptotic for the the scattering phase. Moreover, Weyl asymptotic can also be obtained by representation of the derivative of the spectral shift function involving the trace of the cut-off resolvent (see Robert [36], Bruneau-Petkov [7] and Nakamura [28]).

Concerning the Breit-Wigner approximation for the derivative of the spectral shift function in the Schrödinger case, same kind of results have been obtained in a particular semi-classical set-up by C.Gérard-Martinez-Robert [16] for short range potentials on $\mathbb{R}^n$ and by Petkov-Zworski [32] for a general compactly supported perturbation (see also [7], [6]).

For Dirac operators, Bruneau-Robert [10] established an asymptotic expansion of the scattering phase $s(\lambda)$ and their derivatives in the high energy regime and in the semi-classical regime for λ in a non-trapping energy interval. For an interval $I \subset]-mc^2, mc^2[$ with non critical extremities, Helffer and Robert in [19] gave an asymptotics of the number of the eigenvalues in I for scalar potentials. Nevertheless, we are not aware of works which deal with the link between the derivative of the SSF and resonances for the semi-classical Dirac operators (in the spirit of Petkov-Zworski [32] and Bruneau-Petkov [8]), neither of papers giving the Weyl asymptotic of the spectral shift function for Dirac operators in any interval I .

The purpose of this work is to extend the definition of resonance for analytic perturbations outside a compact set. We define the resonances for the semi-classical Dirac operator as the discrete eigenvalues of the non-selfadjoint operator obtained from the Dirac operator H by a general class of complex distortions of $\mathbb{R}^3$. We prove that the resonances are independent of the distortion (see Section 4). We establish an upper bound for the number of resonances in a compact domain Ω (see Section 5). The second goal of this work is to obtain a meromorphic continuation of the derivative of the spectral shift function $\xi(\lambda, h)$ related to the resonances for the semi-classical Dirac operator (see Section 6). This last is closely related to trace formulae (see [8], [38], [39], [31], [32], [40]) and to resonance expansions (see [42], [11]). Thereafter, in the case where the potential is an electro-magnetic potential, we deduce a Weyl-type asymptotic of the spectral shift function (see Section 7). As a by-product, we obtain an upper bound $O(h^{-2})$ for the number of resonances close to non-critical energy levels in domains of width h (see Subsection 8.1), as well as a Breit-Wigner approximation for the derivative of the SSF (see Subsection 8.2).

2. STATEMENT OF THE RESULTS

We consider the selfadjoint Dirac operator

$$(1) \quad H_0 = -ich \sum_{j=1}^3 \alpha_j \frac{\partial}{\partial x_j} + \beta mc^2,$$

with domain $D(H_0) = H^1(\mathbb{R}^3) \otimes \mathbb{C}^4 \subset \mathcal{H} = L^2(\mathbb{R}^3) \otimes \mathbb{C}^4$, where $h \searrow 0$ is the semi-classical parameter, $m > 0$ is the mass of the Dirac particle and c is the speed of light. The quantities

$\alpha_1, \alpha_2, \alpha_3$ and β are 4×4 Dirac matrices satisfying the anti-commutation relations

$$(2) \quad \begin{cases} \alpha_i \alpha_j + \alpha_j \alpha_i = 2\delta_{ij} I_4, & \text{for } i, j = 1, 2, 3, \\ \alpha_i \beta + \beta \alpha_i = 0, & \text{for } i = 1, 2, 3, \end{cases}$$

and $\beta^2 = I_4$. Here I_n is the $n \times n$ identity matrix. For example, we choose the standard (or Dirac-Pauli) representation of these matrices

$$\alpha_i = \begin{pmatrix} 0 & \sigma_i \\ \sigma_i & 0 \end{pmatrix}, \quad \beta = \begin{pmatrix} I_2 & 0 \\ 0 & -I_2 \end{pmatrix},$$

where $(\sigma_j)_{1 \leq j \leq 3}$ are the 2×2 Pauli matrices:

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

Remark 1. *Most calculations with Dirac matrices can be done without referring to a particular representation (see Appendix 1.A [43, Chap. 1]).*

Let $H_1 = H := H_0 + V$, where V is the multiplication operator by a 4×4 -matrix potential V . We suppose that $V \in C^\infty(\mathbb{R}^3)$ and satisfies the following assumption

(**A_V**) : *V is Hermitian on $\mathbb{R}^3$ and has an analytic extension in the sector*

$$(3) \quad C_{\epsilon,0} := \{z \in \mathbb{C}^3, |\operatorname{Im}(z)| \leq \epsilon |\operatorname{Re}(z)|, |\operatorname{Re}(z)| > R_0\}, \text{ for } 0 < \epsilon < 1.$$

Moreover, for $x \in C_{\epsilon,0}$ it satisfies

$$(4) \quad \|V(x)\| = O(\langle x \rangle^{-\delta}), \quad \delta > 0, \quad \langle x \rangle = (1 + |x|^2)^{\frac{1}{2}}.$$

The free Dirac operator H_0 has only essential spectrum $\sigma_{ess}(H_0) =]-\infty, -mc^2] \cup [mc^2, +\infty[$. Under the assumption (**A_V**) the operator H_1 is a selfadjoint operator. Using Weyl theorem, we have $\sigma_{ess}(H_1) = \sigma_{ess}(H_0)$.

For $\theta \in D_\epsilon := \{\theta \in \mathbb{C}, |\theta| \leq r_\epsilon := \frac{\epsilon}{\sqrt{1+\epsilon^2}}\}$, we denote

$$H_{1,\theta} = H_\theta := U_\theta H_0 U_\theta^{-1} + U_\theta V U_\theta^{-1} = H_{0,\theta} + U_\theta V U_\theta^{-1},$$

where U_θ is the one-parameter family of distortions defined below (see Section 3).

For θ_0 fixed in $D_\epsilon^+ := D_\epsilon \cap \{\theta \in \mathbb{C}, \operatorname{Im}(\theta) \geq 0\}$, we define

$$\Gamma_{\theta_0} := \{\pm c \sqrt{\frac{\lambda}{(1+\theta_0)^2} + m^2 c^2} \in \mathbb{C}, \lambda \in [0, +\infty[\},$$

and

$$S_{\theta_0} := \{z \in \bigcup_{\theta \in D_\epsilon^+} \Gamma_\theta; \arg(1+\theta) < \arg(1+\theta_0), \frac{1}{|1+\theta|} < \frac{1}{|1+\theta_0|}\}.$$

The square root $\sqrt{z}$ is defined such that for $z \in \mathbb{C} \setminus]-\infty, 0]$, $\operatorname{Re}(\sqrt{z}) > 0$.

For $\theta \in D_\epsilon^+$, $\arg(1+\theta_0) \leq \arg(1+\theta)$, $\frac{1}{|1+\theta_0|} \leq \frac{1}{|1+\theta|}$, we prove that the spectrum of H_θ is discrete in S_θ and independent of θ in S_{θ_0} . This justifies the following definition

Definition 1. *The resonances of H in $S_{\theta_0} \cup \mathbb{R}$ are the eigenvalues of H_{θ_0} . The multiplicity of a resonance z is the geometric multiplicity of z considered as an eigenvalue of H_{θ_0} . We will denote $\operatorname{Res}(H)$ the set of resonances.*

The most important advantage of this definition is that the resonances can be computed by solving a non-selfadjoint eigenvalue problem.

Remark 2. *The resonances of H in $\{z \in \mathbb{C}; \operatorname{Re}(z) \in]-mc^2, mc^2[\}$ are the real eigenvalues of H .*




Fig. The set S_{θ_0}

Now, we would like to find symmetry properties which makes it possible to limit our study of the resonances in a domain Ω which satisfies $(\mathbf{A}_{\Omega}^+)$, with

$(\mathbf{A}_{\Omega}^{\pm})$: Ω is an open simply connected and relatively compact subset of $\{z \in \mathbb{C}; \pm \operatorname{Re}(z) > mc^2\}$ such that $\Omega \cap \{\pm \operatorname{Im}(z) > 0\} \neq \emptyset$ and there exists $\theta_0 \in D_{\epsilon}^+$ such that $\overline{\Omega} \cap \Gamma_{\theta_0} = \emptyset$.

Proposition 1. *Let H^- be the selfadjoint Hamiltonian*

$$H^- = H_0 - U_c \overline{V}(x) U_c^{-1},$$

where $U_c = i\beta\alpha_2$ is an unitary matrix 4×4 and $\overline{V}$ is the conjugate of V . Then the following assertions are equivalents:

- (i) *The complex value z is a resonance of H .*
- (ii) *The symmetric of the conjugate $-\bar{z}$ is a complex eigenvalue of $U_{\bar{\theta}} H^- U_{\bar{\theta}}^{-1}$.*

Moreover, the multiplicity of z is equal to the multiplicity of $-\bar{z}$ considered as an eigenvalue of $U_{\bar{\theta}} H^- U_{\bar{\theta}}^{-1}$.

Proposition 2. *Let $\overline{H}$ be the selfadjoint Hamiltonian*

$$\overline{H} = -ic \sum_{j=1}^3 \alpha'_j \partial_{x_j} + \beta' mc^2 + \overline{V},$$

where $\alpha'_1 = -\alpha_1$, $\alpha'_2 = \alpha_2$, $\alpha'_3 = -\alpha_3$, $\beta' = \beta$ are matrices which satisfy the anti-commutation relations (2) and $\overline{V}$ is the conjugate of V . Then the following assertions are equivalent:

- (i) *The complex value z is a resonance of H .*
- (ii) *The conjugate $\bar{z}$ is a complex eigenvalue of $U_{\bar{\theta}} \overline{H} U_{\bar{\theta}}^{-1}$.*

Moreover, the multiplicity of z is equal to the multiplicity of $\bar{z}$ considered as an eigenvalue of $U_{\bar{\theta}} \overline{H} U_{\bar{\theta}}^{-1}$.

Using the same type of approach as in [39], we construct an operator $\widehat{H}_{j,\theta}$ for $j = 0, 1$, so that

$$\widehat{H}_{j,\theta} - H_{j,\theta} = K_j = O(1), \text{ has finite rank } O(h^{-3}),$$

and $\|(\widehat{H}_{j,\theta} - z)^{-1}\| = O(1)$, uniformly for $z \in \overline{\Omega}$ (see Subsection 5.1).

Using this construction we establish an upper bound of the number of resonances:

Theorem 1. (Upper bound) *Assume that V satisfies the assumption $(\mathbf{A}_V)$ with $\delta > 0$. Let Ω be a complex domain satisfying the assumption $(\mathbf{A}_{\Omega}^{\pm})$, then*

$$\#\text{Res}(H) \cap \Omega \leq C(\Omega)h^{-3}.$$

For a pair of self-adjoint operators $(H_0, H_0 + V)$ where V satisfies the assumption $(\mathbf{A}_V)$ with $\delta > 3$, (see [10], [36], [35]), the spectral shift function $\xi(\lambda, h)$ is a distribution in $\mathcal{D}'(\mathbb{R})$ such that its derivative satisfies:

$$(5) \quad \langle \xi'(\lambda, h), f(\lambda) \rangle_{\mathcal{D}'(\mathbb{R}), \mathcal{D}(\mathbb{R})} = \text{tr}(f(H_1) - f(H_0)), \quad f(\lambda) \in C_0^\infty(\mathbb{R}).$$

By the Birman-Krein theory, the SSF is in $L_{loc}^1(\mathbb{R})$ and coincides with the scattering phase (see [44]).

Our principal result is a meromorphic continuation of the derivative of the spectral shift function $\xi(\lambda, h)$.

Theorem 2. (Representation formula) *Assume that V satisfies the assumption $(\mathbf{A}_V)$ with $\delta > 3$. Let Ω be a complex domain satisfying the assumption $(\mathbf{A}_{\Omega}^{\pm})$ and $W \Subset \Omega$ be an open simply connected set which is symmetric with respect to the real axis. Assume that $I = W \cap \mathbb{R}$ is an interval. Then for all $\lambda \in I$ we have the representation:*

$$(6) \quad \xi'(\lambda, h) = \frac{1}{\pi} \text{Im } r(\lambda, h) + \sum_{\substack{w \in \text{Res}(H_1) \cap \Omega \\ \text{Im } w \neq 0}} \frac{-\text{Im } w}{\pi |\lambda - w|^2} + \sum_{w \in \text{Res}(H_1) \cap I} \delta_w(\lambda),$$

where $r(z, h) = g(z, h) - \overline{g(\bar{z}, h)}$, $g(z, h)$ is a holomorphic function in Ω which satisfies the following estimate:

$$(7) \quad |g(z, h)| \leq C(W)h^{-3}, \quad z \in W,$$

with $C(W) > 0$ independent of $h \in]0, h_0]$. Here $\delta_w(\cdot)$ is the Dirac mass at $w \in \mathbb{R}$.

Remark 3. *This theorem can be extended to the pair operators $(H_0 + V_1, H_0 + V_2)$, where V_1, V_2 are two 4×4 Hermitian potential matrices satisfying $(\mathbf{A}_V)$ with $\delta > 0$ and $V = V_2 - V_1$ satisfies the assumption $(\mathbf{A}_V)$ with $\delta > 3$ (see Theorem 5).*

As a corollary of the last theorem, we have a Sjöstrand type local trace formula (see Theorem 6).

Now we discuss a Weyl type asymptotic of the spectral shift function $\xi(\lambda, h)$ in the case where V is an electro-magnetic potential

$$(8) \quad V(x) = e(-\alpha \cdot A + v)(x) = - \sum_{j=1}^3 \alpha_j \cdot eA_j(x) + ev(x),$$

satisfying the assumption $(\mathbf{A}_V)$ with $\delta > 3$. Here $e < 0$ is the charge of the Dirac particle. We assume that, the electric potential $v(x) = \begin{pmatrix} v_+(x)I_2 & 0 \\ 0 & v_-(x)I_2 \end{pmatrix}$ where v_+, v_- are C^∞ scalar functions satisfying

$$(9) \quad |e(v_+ - v_-)(x)| < 2mc^2,$$

and $A = (A_1, A_2, A_3)$ is a magnetic vector potential where A_1, A_2, A_3 are C^∞ scalar functions.

For any $(x, \xi) \in \mathbb{R}^6$, the semi-classical symbols of H_ν , $\nu = 0, 1$ are the matrices

$$(10) \quad \mathcal{D}_\nu(x, \xi) = \alpha \cdot (c\xi - \nu eA(x)) + \beta mc^2 + \nu ev(x), \quad \alpha = (\alpha_1, \alpha_2, \alpha_3),$$

which is Hermitian and has two eigenvalues

$$(11) \quad H_\nu^\pm(x, \xi) = \pm \left(|c\xi - \nu eA(x)|^2 + \left(mc^2 + \nu \frac{e}{2}(v_+ - v_-) \right)^2 \right)^{\frac{1}{2}} + \nu \frac{e}{2}(v_+ + v_-),$$

of multiplicity two. The function $H_1^+(x, \xi)$ is the Hamiltonian for a relativistic classical particle and the other $H_1^-(x, \xi)$ can be considered as the one for its anti-particle (see [14], [43], [46]). Moreover, from (9) the two Hamiltonians $H_1^\pm(x, \xi)$ are smooth functions.

For $\nu = 0, 1$, the matrix

$$(12) \quad \Pi_\nu^\pm(x, \xi) = \frac{1}{2} \left(1 + \frac{\mathcal{D}_\nu(x, \xi) - \nu ev(x) + \nu \frac{e}{2}(v_+ - v_-)\beta}{H_\nu^\pm(x, \xi) - \nu \frac{e}{2}(v_+ + v_-)} \right),$$

is the orthogonal projection on the eigenspace $\mathcal{E}_\nu^\pm(x, \xi)$ of $\mathcal{D}_\nu(x, \xi)$ corresponding to the eigenvalue $H_\nu^\pm(x, \xi)$.

Definition 2. *A real λ is a noncritical energy level for H_1 if for all $(x, \xi) \in \mathbb{R}^6$, with $H_1^\pm(x, \xi) = \lambda$, we have $\nabla_{x, \xi} H_1^\pm(x, \xi) \neq 0$.*

Theorem 3. (Weyl formula) *Assume that the potential V is an electro-magnetic potential given by (8) and satisfying the assumption $(\mathbf{A}_V)$ with $\delta > 3$. For all λ, λ_1 noncritical energy levels for H_1 such that $\pm mc^2 \notin]\lambda_1, \lambda[$ and $h \in]0, h_0[$, we have the asymptotic expansion*

$$(13) \quad \xi(\lambda, h) - \xi(\lambda_1, h) = w(\lambda, \lambda_1)h^{-3} + O(h^{-2}).$$

Here the $O(h^{-2})$ is uniform for λ (resp. λ_1) in a small interval I_2 (resp. I_1). The first term $w(\lambda, \lambda_1) \in C^\infty(I_2 \times I_1)$ is given by

$$w(\lambda, \lambda_1) = w(\lambda) - w(\lambda_1),$$

with:

$$(14) \quad w(\lambda) = \frac{1}{3\pi^2} \int_{\mathbb{R}^3} W_+(\lambda, v_+, v_-) - W_+(\lambda, 0, 0) - W_-(\lambda, v_+, v_-) + W_-(\lambda, 0, 0) \, dx$$

$$\text{where } W_{\pm}(\lambda, a, b) = \left(\left(\lambda - \frac{e(a+b)}{2} \right)_{\pm}^2 - \left(mc^2 + \frac{e(a-b)}{2} \right)^2 \right)_{+}^{\frac{3}{2}}.$$

The Theorem 3 can be extended to the pair operators $(H_0 + V_1, H_0 + V_2)$ (see Remark 10).

Remark 4. *The two formulae (13) and (14) give in particular a Weyl type asymptotic of the counting function of the number of eigenvalues of H_1 between two values in the interval $] -mc^2, mc^2[$. In the case of scalar potential $v(v_+ = v_-)$, this result was proved by Helffer-Robert [19] without the analytic assumption at infinity.*

To prove Theorem 3 we construct, in Appendix A (see Theorem 9), a parametrix at small time of the propagator of the Dirac equation in an external electro-magnetic field (see also Yajima [46] for a scalar electric potential cases).

As a direct result of the last theorem we deduce an upper bound $O(h^{-2})$ for the number of resonances close to non-critical energy levels in domains of width h (see Proposition 11) and a Breit-Wigner approximation for the derivative of the spectral shift function $\xi(\lambda, h)$ (see Theorem 8).

3. DISTORTION FOR THE FREE DIRAC OPERATOR

In this section, we start with the definition of the deformation for the free Dirac operator by analytic distortion (in the spirit of Hunziker [21]) and we will calculate the essential spectrum for the distorted free Dirac operator. Here, h does not play any role, and can be taken equal to 1. Let us now introduce the one-parameter family of unitary distortion

$$U_{\theta}f(x) = J_{\phi_{\theta}(x)}^{\frac{1}{2}}f(\phi_{\theta}(x)), \quad \theta \in \mathbb{R}, \quad f \in (S(\mathbb{R}^3))^4,$$

where $\phi_{\theta}(x) = x + \theta g(x)$ and $g : \mathbb{R}^3 \mapsto \mathbb{R}^3$ is a smooth function. Let $J_{\phi_{\theta}(x)} = \det(I + \theta \nabla g(x))$ be the Jacobian of $\phi_{\theta}(x)$.

We suppose that g satisfies the assumption

$$(\mathbf{A}_g) \left\{ \begin{array}{l} \text{(i) } \sup_{x \in \mathbb{R}^3} \|\nabla g(x)\| = M^{-1} < +\infty. \\ \text{(ii) } g(x) = 0, \text{ in the compact } B(0, R_0), \text{ (see (3)).} \\ \text{(iii) } g(x) = x, \text{ outside of a compact set } K(\supset B(0, R_0)). \end{array} \right.$$

Lemma 1. *For $\theta \in]-M, M[$, U_{θ} can be extended as an unitary operator on $\mathcal{H}$.*

Proof. Since $|\theta| < M$, we have $\|\theta \nabla g(x)\| < 1$, and

$$(\nabla \phi_{\theta}(x))^{-1} = (I + \theta \nabla g(x))^{-1} = \sum_{n=0}^{\infty} (-1)^n (\theta)^n (\nabla g(x))^n.$$

The function $\phi_\theta(x)$ is injective and $\phi_\theta(\mathbb{R}^3) = \mathbb{R}^3$, consequently $\phi_\theta(x)$ is a diffeomorphism from $\mathbb{R}^3$ to $\mathbb{R}^3$. The inverse of U_θ is given by

$$U_\theta^{-1}u = J_{\phi_\theta(x)}^{-\frac{1}{2}}u(\phi_\theta^{-1}(x)) : (L^2(\mathbb{R}^3))^4 \mapsto (L^2(\mathbb{R}^3))^4.$$

The lemma follows from the relations

$$U_\theta U_\theta^{-1} = U_\theta^{-1} U_\theta = I_{(L^2(\mathbb{R}^3))^4} \text{ and } \|U_\theta f\|_{\mathcal{H}} = \|U_\theta^{-1} f\|_{\mathcal{H}} = \|f\|, \quad \forall f \in (L^2(\mathbb{R}^3))^4.$$

□

Definition 3. We denote by $\mathcal{B}$, the space of functions $f = (f_i)_{1 \leq i \leq 4}$ such that $f_i(x)$ has an analytic continuation in $C_{\epsilon,0}$ and $\lim_{\substack{|z| \rightarrow \infty \\ z \in C_{\epsilon,0}}} |z|^k f_i(z) = 0$, for all $k \in \mathbb{N}$ and $\epsilon \in]0, 1[$ (see (3)).

Lemma 2. The subspace $\mathcal{B}$ is dense in $\mathcal{H}$.

Proof. The subspace $\mathcal{B}$ contains vectors of Hermite functions and the linear combinations of Hermite functions are dense in $L^2(\mathbb{R}^3)$. □

Proposition 3. Let be $D_{\epsilon,M} = D_\epsilon \cap \{\theta \in \mathbb{C}; |\theta| < M\}$. We have the two assertions:

- (i) For all $f \in \mathcal{B}$, $\theta \in D_{\epsilon,M} \mapsto U_\theta f$ is analytic.
- (ii) For all $\theta \in D_{\epsilon,M}$, $U_\theta \mathcal{B}$ is dense in $\mathcal{H}$.

Proof. In the order to prove (i), we show that $\theta \mapsto \langle U_\theta f, g \rangle$ is analytic for all $g \in \mathcal{H}$. Let $R \gg 1$ be such that $K \subset B(0, R) = \{x, |x| < R\}$.

- In $\{x, |x| < R\}$: since $f \in \mathcal{B}$, then $\theta \mapsto \int_{|x| < R} J_{\phi_\theta}^{\frac{1}{2}} f(\phi_\theta(x)) \overline{g(x)} dx$, is clearly analytic for all $g \in \mathcal{H}$.
- In $\{x, |x| > R\}$: we have $g(x) = x$, consequently $\phi_\theta(x) = (1 + \theta)x$. We remark that

$$|\operatorname{Im}(\phi_\theta(x))| = |\operatorname{Im}(\theta x)| \leq |\operatorname{Im}(\theta)| |x| = \frac{|\operatorname{Im}(\theta)| |\operatorname{Re}(\phi_\theta(x))|}{|1 + \operatorname{Re}(\theta)|}.$$

If $|\theta| \leq r_\epsilon = \frac{\epsilon}{\sqrt{1+\epsilon^2}}$, $0 < \epsilon < 1$, then

$$|\operatorname{Im}(\phi_\theta(x))| \leq \epsilon |\operatorname{Re}(\phi_\theta(x))|.$$

According to the definition of $\mathcal{B}$ we have

$$|f(\phi_\theta(x))| \leq \frac{C_k}{|\phi_\theta(x)|^k} \leq \frac{C_{k,\epsilon}}{|x|^k}, \quad \forall |x| \geq R, \quad \forall \theta \in D_{\epsilon,M}, \quad k \in \mathbb{N}$$

then, $\theta \mapsto \int_{|x| \geq R} J_{\phi_\theta}^{\frac{1}{2}} f(\phi_\theta) \overline{g(x)} dx$ is analytic.

(ii) Let $h(x) \in (C_0^\infty(\mathbb{R}^3))^4$. We denote

$$h_k(x) = \left(\frac{k}{\pi}\right)^{\frac{3}{2}} \int e^{-k(x-y-\theta g(y))^2} h(y) J_{\phi_\theta(y)} dy,$$

which is clearly in $\mathcal{B}$.

Using $\left(\frac{k}{\pi}\right)^{\frac{3}{2}} \int e^{-k(x-y-\theta g(y))^2} J_{\phi_\theta(y)} dy = \left(\frac{k}{\pi}\right)^{\frac{3}{2}} \int e^{-kz^2} dz = 1$, we get

$$h(x) - h_k(\phi_\theta(x)) = \left(\frac{k}{\pi}\right)^{\frac{3}{2}} \int e^{-k(\phi_\theta(x)-\phi_\theta(y))^2} (h(x) - h(y)) J_{\phi_\theta(y)} dy.$$

The last term tends to 0 when $k \rightarrow +\infty$. Consequently, we have

$$h_k \circ \phi_\theta(x) \xrightarrow{k \rightarrow +\infty} h(x), \quad \text{in } \mathcal{H}.$$

□

Remark 5. One can always choose g satisfying the assumption $(\mathbf{A}_g)$ with $M > r_\epsilon = \frac{\epsilon}{\sqrt{1+\epsilon^2}}$. In that case, we have $D_{\epsilon,M} = D_\epsilon$.

Lemma 3. For $\theta \in D_\epsilon$, we have

$$H_{0,\theta} := U_\theta H_0 U_\theta^{-1} = \frac{1}{1+\theta} (-ic \sum_{j=1}^3 \alpha_j \frac{\partial}{\partial x_j}) + \beta mc^2 + Q_\theta(x, \partial_{x_j}),$$

where $Q_\theta(x, \partial_{x_j}) = \sum_{|\alpha| \leq 1} a_\alpha(x, \theta) \partial_{x_j}^\alpha$ is such that:

- (i) $\theta \mapsto a_\alpha(x, \theta)$ is an analytic function bounded by $O(\theta)$.
- (ii) $x \mapsto a_\alpha(x, \theta) \in (C_0^\infty(\mathbb{R}^3))^4$.

In particular $\theta \mapsto H_{0,\theta}$ is an analytic family of type A of domain $D(H_0)$ (see Kato [23], for the definition of an analytic family of type A).

Proof. We denote $\partial_j = \frac{\partial}{\partial x_j}$ and we calculate the term $U_\theta \partial_j U_\theta^{-1}$.

$$\begin{aligned} U_\theta \partial_j U_\theta^{-1} f(x) &= U_\theta \partial_j \left(J_{\phi_\theta}^{-\frac{1}{2}} f(\phi_\theta^{-1}(x)) \right) \\ &= U_\theta \left(\partial_j \left(J_{\phi_\theta}^{-\frac{1}{2}} \right) \cdot f(\phi_\theta^{-1}(x)) + J_{\phi_\theta}^{-\frac{1}{2}} (\partial_j f(\phi_\theta^{-1}(x))) \right) \\ &= -\frac{1}{2} J_{\phi_\theta}^{-1} \partial_j J_{\phi_\theta(x)} f(x) + \sum_{k=1}^3 \partial_k f(x) \left(\partial_j \phi_{\theta,k}^{-1} \right) (\phi_\theta(x)), \end{aligned}$$

with $\phi_\theta^{-1}(x) = (\phi_{\theta,1}^{-1}(x), \phi_{\theta,2}^{-1}(x), \phi_{\theta,3}^{-1}(x))$.

We remark that $\phi_\theta^{-1}(x) = \frac{x}{1+\theta}$ outside of the compact set K , then

$$\sum_{k=1}^3 \partial_k f(x) \left(\partial_j \phi_{\theta,k}^{-1} \right) (\phi_\theta(x)) = \frac{1}{1+\theta} \partial_j f(x), \text{ outside of } K.$$

Let $\chi_K \in C_0^\infty(\mathbb{R}^3)$, $0 \leq \chi_K \leq 1$, be equal to 1 on K and 0 outside a compact set which contains K . We have

$$U_\theta \partial_j U_\theta^{-1} f(x) = -\frac{1}{2} J_{\phi_\theta}^{-1} \partial_j J_{\phi_\theta(x)} f(x) + \frac{1}{1+\theta} \partial_j f(x) (1 - \chi_K) + \sum_{k=1}^3 \partial_k f(x) \left(\partial_j \phi_{\theta,k}^{-1} \right) (\phi_\theta(x)) \chi_K.$$

Since $\partial_j J_{\phi_\theta(x)}$ has a compact support, then

$$(15) \quad U_\theta \partial_j U_\theta^{-1} = \frac{1}{1+\theta} \partial_j + q_\theta(x, \partial_{x_j})$$

with $q_\theta(x, \partial_{x_j})$ satisfying the hypothesis of $Q_\theta(x, \partial_{x_j})$.

Now we just have to multiply (15) by $-ic\alpha_j$, to sum on all values of j and add βmc^2 to both hands. The estimate $a_\alpha(x, \theta) = O(\theta)$ is clear using that $Q_0(x, \partial_{x_j}) = 0$ and the analytic property of $\theta \mapsto a_\alpha(\cdot, \theta)$. □

Lemma 4. *Let $P_\theta = \frac{1}{1+\theta}(-ic \sum_j \alpha_j \frac{\partial}{\partial x_j}) + \beta mc^2$. Then*

$$\sigma(P_\theta) = \sigma_{ess}(P_\theta) = \Gamma_\theta = \{z \in \mathbb{C}; z = \pm c(\frac{\lambda}{(1+\theta)^2} + m^2 c^2)^{\frac{1}{2}}, \lambda \in [0, +\infty[\}.$$

Proof. Let $\mathcal{F}$ be the Fourier transform and

$$\begin{aligned} K(\theta) = \mathcal{F} P_\theta \mathcal{F}^{-1} &= \frac{c}{1+\theta} \sum_j \alpha_j \xi_j + mc^2 \beta \\ &= \begin{pmatrix} mc^2 I_2 & (\frac{c}{1+\theta})(\sigma_1 \xi_1 + \sigma_2 \xi_2 + \sigma_3 \xi_3) \\ (\frac{c}{1+\theta})(\sigma_1 \xi_1 + \sigma_2 \xi_2 + \sigma_3 \xi_3) & -mc^2 I_2 \end{pmatrix} \end{aligned}$$

where $\xi = (\xi_1, \xi_2, \xi_3) \in \mathbb{R}^3$ and $\alpha_j \xi_j$ is the multiplication operator by the 4×4 matrix $\alpha_j \xi_j$. Here $\sigma_1, \sigma_2, \sigma_3$ are the 2×2 Pauli matrices. The spectrum of P_θ coincide with the spectrum of the multiplication operator $K(\theta)$. We easily prove that

$$\sigma(K(\theta)) = \sigma_{ess}(K(\theta)) = \Gamma_\theta = \{z \in \mathbb{C}; z = \pm c(\frac{\lambda}{(1+\theta)^2} + m^2 c^2)^{\frac{1}{2}}, \lambda \in [0, +\infty[\},$$

and we deduce the lemma. $\square$

The principal branch of the square root function is holomorphic on the set $\mathbb{C} \setminus]-\infty, 0]$. Let $S_{D_\epsilon} = \{z = \frac{\lambda}{(1+\theta)^2} + m^2 c^2, \theta \in D_\epsilon, \lambda \in [0, +\infty[\}$. Since,

$$S_{D_\epsilon} \subset]0, +\infty[e^{-\frac{\pi}{2}, \frac{\pi}{2}}[,$$

the square root $z \mapsto z^{\frac{1}{2}}$ is holomorphic on S_{D_ϵ} .

Lemma 5. *For $H_{0,\theta}, P_\theta$, defined as above, we have $\sigma_{ess}(H_{0,\theta}) = \sigma_{ess}(P_\theta)$.*

Proof. We want to use Kato's theorem [23, Th.4.5.35]. For $\lambda \gg 1$, $\lambda \in \mathbb{R}$ and Q_θ defined in Lemma 3, we have

$$(H_{0,\theta} - i\lambda) = (1 + Q_\theta(P_\theta - i\lambda)^{-1})(P_\theta - i\lambda).$$

Since $(P_\theta - i\lambda)^{-1} \in \mathcal{L}(\mathcal{H}, (H^1)^4)$ and $Q_\theta(P_\theta - i\lambda)^{-1} = O(\frac{\theta}{\lambda})$, we obtain that $i\lambda \in \rho(H_{0,\theta}) = \mathbb{C} \setminus \sigma(H_{0,\theta})$. To apply the Kato Theorem, it is enough to show that

$$(16) \quad (H_{0,\theta} - i\lambda)^{-1} - (P_\theta - i\lambda)^{-1} \text{ is compact.}$$

Using the resolvent equation, we have

$$(H_{0,\theta} - i\lambda)^{-1} - (P_\theta - i\lambda)^{-1} = (H_{0,\theta} - i\lambda)^{-1} Q_\theta (P_\theta - i\lambda)^{-1}.$$

with $Q_\theta(x, \partial_{x_j}) = \sum_{|\alpha| \leq 1} a_\alpha(x, \theta) \partial_{x_j}^\alpha$ compactly supported. Since the operator $(H_{0,\theta} - i\lambda)^{-1} Q_\theta$ is bounded and $\mathbf{1}_{\text{supp}(Q_\theta)}(P_\theta - i\lambda)^{-1}$ is compact, the assertion (16) holds. $\square$

4. DEFINITION OF RESONANCES

In this section we distort the perturbed Dirac operator $H = H_0 + V$, where the potential V satisfies the assumption **(A_V)** and we define the resonances for the semi-classical Dirac operator.

The distorted Dirac operator is denoted by

$$H_\theta = U_\theta H_0 U_\theta^{-1} + U_\theta V U_\theta^{-1} = H_{0,\theta} + V(\phi_\theta(x)).$$

Proposition 4. *We suppose that the potential V satisfies the assumption **(A_V)**, then*

(i) $\theta \in D_\epsilon \mapsto H_\theta = H_{0,\theta} + V(\phi_\theta(x))$ is an analytic family of type A.

$$(ii) \quad \sigma_{ess}(H_\theta) = \Gamma_\theta.$$

Proof. The assertion (i) is clear since $H_{0,\theta}$ is an analytic family of type A and V satisfies the assumption **(A_V)**.

Now, we prove (ii) as in the proof of Lemma 5. For $\lambda \gg 1$, $i\lambda \in \rho(H_\theta)$ and

$$(17) \quad (H_\theta - i\lambda)^{-1} - (H_{0,\theta} - i\lambda)^{-1} = (H_\theta - i\lambda)^{-1} V(\phi_\theta(x)) (H_{0,\theta} - i\lambda)^{-1}.$$

Since the operator $V(\phi_\theta(x))(H_{0,\theta} - i\lambda)^{-1}$ is compact (see **(A_V)**) and the resolvent $(H_\theta - i\lambda)^{-1}$ is bounded, the difference $(H_\theta - i\lambda)^{-1} - (H_{0,\theta} - i\lambda)^{-1}$ is compact. According to Kato's theorem [23, Theorem.4.5.35] and to the lemmas 4, 5, we obtain (ii). $\square$

We denote

$$\Sigma = \{z \in \mathbb{C}; \operatorname{Im}(z) \geq 0, \operatorname{Re}(z) > -mc^2\} \cup \{z \in \mathbb{C}; \operatorname{Im}(z) \leq 0, \operatorname{Re}(z) < mc^2\} \setminus \sigma(H).$$

Theorem 4. *With the notations used above and taking $\theta_0 \in D_\epsilon^+ = D_\epsilon \cap \{\operatorname{Im}(\theta) \geq 0\}$, we have:*

- (i) *For all $f, g \in \mathcal{B}$, the function: $z \in \Sigma \mapsto M_{f,g}(z) = \langle (z - H)^{-1} f, g \rangle$ has a meromorphic extension on S_{θ_0} .*
- (ii) *The poles of $M_{f,g}(z)$ are the eigenvalues of H_{θ_0} .*
- (iii) *These poles are independent of the family U_{θ_0} .*
- (iv) *$\sigma_d(H_{\theta_0}) \cap \Sigma = \emptyset$, where $\sigma_d(H_{\theta_0})$ is the discrete spectrum of the operator H_{θ_0} .*

Proof. (i) Since U_θ is unitary for $\theta \in \mathbb{R}$,

$$M_{f,g}(z) = \langle (z - H)^{-1} f, g \rangle = \langle (z - H_\theta)^{-1} U_\theta f, U_\theta g \rangle.$$

We denote

$$(18) \quad M_{f,g,\theta}(z) = \langle (z - H_\theta)^{-1} U_\theta f, U_\theta g \rangle, \quad \text{for } \theta \in D_\epsilon.$$

According to (i) of Proposition 4 and to the definition of U_θ , the functions $\theta \mapsto (z - H_\theta)^{-1}$, $\theta \mapsto U_\theta f$ and $\theta \mapsto \langle \psi, U_\theta g \rangle$ are analytic on D_ϵ for all $\psi \in \mathcal{H}$ and any $z \in \Sigma$.

Thus, for $z \in \Sigma$, the function $\theta \mapsto M_{f,g,\theta}(z)$ is analytic on D_ϵ . Since $M_{f,g,\theta}(z)$ is independent of θ on the real axis and according to the uniqueness of the prolongation, its extension is independent of θ .

Now, we fix $\theta_0 \in D_\epsilon^+$. Since $S_{\theta_0} \cap \sigma_{ess}(H_{\theta_0}) = \emptyset$, the function $z \in \Sigma \mapsto M_{f,g,\theta_0}(z)$ has a meromorphic extension in S_{θ_0} .

(ii) First, let $z \in S_{\theta_0}$ be a pole of $M_{f,g}(z)$ which is equal to $M_{f,g,\theta_0}(z)$ for $\theta_0 \in D_\epsilon^+$. Then $z \in \sigma_d(H_{\theta_0}) \cap S_{\theta_0}$ (see proof of (i)).

Now, let $e \in \sigma_d(H_{\theta_0}) \cap S_{\theta_0}$. There exists $u \in \mathcal{H}$ such that $\|u\| = 1$ and $H_{\theta_0} u = eu$. Let γ be a small disk centered at e such that $\overset{\circ}{\gamma} \cap \sigma(H_{\theta_0}) = \{e\}$ and Γ be the positively oriented boundary of γ .

Let us introduce the projector

$$\Pi = \frac{1}{2i\pi} \int_{\Gamma} (z - H_{\theta_0})^{-1} dz; \quad \Pi u = u.$$

Since $\overline{U_{\theta_0} \mathcal{B}} = \mathcal{H} = \overline{U_{\bar{\theta}_0} \mathcal{B}}$ (see Proposition 3), there exist $f_n, g_n \in \mathcal{B}$ such that

$$|u - U_{\theta_0} f_n| \leq \frac{1}{n} \quad \text{and} \quad |u - U_{\bar{\theta}_0} g_n| \leq \frac{1}{n}, \quad n \in \mathbb{N}.$$

Therefore, as n goes to infinity, we have

$$\begin{aligned}
\frac{1}{2i\pi} \int_{\Gamma} \langle (z - H_{\theta_0})^{-1} U_{\theta_0} f_n, U_{\bar{\theta}_0} g_n \rangle dz &= \frac{1}{2i\pi} \int_{\Gamma} \langle (z - H_{\theta_0})^{-1} u, u \rangle dz + o(1) \\
&= \langle \Pi u, u \rangle + o(1) \\
&= \|u\|^2 + o(1) \\
&= 1 + o(1),
\end{aligned}$$

and then,

$$\frac{1}{2i\pi} \int_{\Gamma} \langle (z - H)^{-1} f_n, g_n \rangle dz = 1 + o(1).$$

So that $M_{f_n, g_n}(z)$ admits e as a pole in γ .

The assertion (iii) results from (ii) because $M_{f, g}(z)$ is independent of U_{θ} .

(iv) If there exists $z \in \sigma_d(H_{\theta_0}) \cap \Sigma$, then z is a pole of $\langle (z - H)^{-1} f, g \rangle$, for $f, g \in \mathcal{B}$, but $\langle (z - H)^{-1} f, g \rangle$ is analytic on this domain. We conclude that such z does not exist. $\square$

Remark 6. (i) *It results from (ii) of Theorem 4 that for all $\theta \in D_{\epsilon}^+$, the discrete spectrum $\sigma_d(H)$ is a subset of $\sigma_d(H_{\theta})$.*

(ii) *The previous theorem justifies the definition of the resonances (Definition 1) and using Lemma 4, H_0 has no resonances.*

Remark 7. *If $\theta \in D_{\epsilon}$, then its conjugate $\bar{\theta} \in D_{\epsilon}$. Repeating arguments of the proof of Theorem 4, we have*

- (i) *The function $\theta \mapsto M_{f, g, \bar{\theta}}(z)$ has a analytic extension for $\theta \in D_{\epsilon}$.*
- (ii) *The function $z \in \bar{\Sigma} \mapsto M_{f, g, \bar{\theta}}(z)$ has a meromorphic extension on $S_{\bar{\theta}_0}$,*

where

$$\bar{\Sigma} = \{z \in \mathbb{C}; \operatorname{Im}(z) \geq 0, \operatorname{Re}(z) < mc^2\} \cup \{z \in \mathbb{C}; \operatorname{Im}(z) \leq 0, \operatorname{Re}(z) > -mc^2\} \setminus \sigma(H),$$

and $S_{\bar{\theta}_0}$ is the symmetric of S_{θ_0} with respect to the real axis

$$S_{\bar{\theta}_0} = \{z \in \bigcup_{\theta \in D_{\epsilon}^+} \Gamma_{\bar{\theta}}; \arg(1 + \theta) < \arg(1 + \theta_0), \frac{1}{|1 + \theta|} < \frac{1}{|1 + \theta_0|}\}.$$

Consequently, we obtain (see Theorem 4):

- 1) *The poles of $M_{f, g}(z)$ in $S_{\bar{\theta}_0}$ are the eigenvalues of $H_{\bar{\theta}_0}$.*
- 2) *These poles are independent of the family $U_{\bar{\theta}_0}$.*
- 3) *$\sigma_d(H_{\bar{\theta}_0}) \cap \bar{\Sigma} = \emptyset$.*

The assertions 3) and (ii) prove that the operator $H_{\bar{\theta}}$ has only discrete spectrum in $S_{\bar{\theta}_0}$.

Proof of the Proposition 1. We consider the anti-linear application on $\mathcal{H}$,

$$C : \psi \mapsto U_c \bar{\psi} = i\beta\alpha_2 \bar{\psi}.$$

Then, we have

$$\begin{aligned}
C H_{\theta} C^{-1} &= -H_{0, \bar{\theta}} + CV \circ \phi_{\theta}(x) C^{-1} \\
&= -\left(H_{0, \bar{\theta}} - U_c \overline{V \circ \phi_{\theta}(x)} U_c^{-1}\right).
\end{aligned}$$

Using that V is analytic, we get $\overline{V \circ \phi_\theta(x)} = \overline{V} \circ \phi_{\bar{\theta}}(x)$. Then,

$$\begin{aligned} C H_\theta C^{-1} &= - (H_{0,\bar{\theta}} - U_c \overline{V} \circ \phi_{\bar{\theta}}(x) U_c^{-1}) \\ &= - (H_{0,\bar{\theta}} - U_c U_{\bar{\theta}} \overline{V}(x) U_{\bar{\theta}}^{-1} U_c^{-1}). \end{aligned}$$

We recall that $U_\theta f(x) = J_{\phi_\theta(x)}^{\frac{1}{2}} f(\phi_\theta(x))$. Since $U_c U_\theta = U_\theta U_c$, we obtain

$$\begin{aligned} C H_\theta C^{-1} &= - (H_{0,\bar{\theta}} - U_{\bar{\theta}} U_c \overline{V}(x) U_c^{-1} U_{\bar{\theta}}^{-1}) \\ &= - U_{\bar{\theta}} (H_0 - U_c \overline{V}(x) U_c^{-1}) U_{\bar{\theta}}^{-1} = - H_{\bar{\theta}}^-. \end{aligned}$$

Consequently, $C (H_\theta - z) C^{-1} = -(H_{\bar{\theta}}^- + \bar{z})$, and the property follows. $\square$

Proof of the Proposition 2.

By definition of H_θ , we have

$$(19) \quad \overline{H_\theta - z} = U_{\bar{\theta}} (\overline{H_0} + \overline{V} - \bar{z}) U_{\bar{\theta}}^{-1}.$$

Using that $\bar{\alpha}_1 = \alpha_1$, $\bar{\alpha}_2 = -\alpha_2$, $\bar{\alpha}_3 = \alpha_3$, $\bar{\beta} = \beta$, we find

$$\overline{H_0} = ic \sum_{j=1}^3 \bar{\alpha}_j \frac{\partial}{\partial x_j} + \beta m c^2 = -ic \sum_{j=1}^3 \alpha'_j \frac{\partial}{\partial x_j} + \beta' m c^2$$

and

$$\overline{H_0} + \overline{V} = \overline{H}.$$

Using the last relation and the equation (19), we obtain Proposition 2. $\square$

Finally, the study of resonances in a domain of the complex plan $\mathbb{C}$ is reduced to study the resonances in $\Omega \cap \{z \in \mathbb{C}, \text{Im}(z) < 0\}$, with Ω satisfying the assumption $(\mathbf{A}_\Omega^+)$ (see Fig.1).

5. UPPER BOUND FOR THE NUMBER OF RESONANCES

In this section, we establish an upper bound of the number of resonances in a compact domain Ω . For this purpose we construct an operator $\widehat{H}_\theta : D(H_0) \rightarrow \mathcal{H}$ with some properties (see Proposition 6). According to the Section 4, it is sufficient to treat the case where Ω satisfies the assumption $(\mathbf{A}_\Omega^+)$.

We shall use the theory of h -pseudo-differential operator (see [12], [34]). Let m be an order function on $\mathbb{R}^{2n}$ (i.e. there are $C_0, N_0 > 0$, such that $m(x) \leq C_0 \langle x - y \rangle^{N_0} m(y)$). The space $\mathcal{S}^p(m)$ is the set of $a(x, \xi; h) \in C^\infty(\mathbb{R}^{2n}) \otimes \mathbb{C}^4$ such that for every $\alpha \in \mathbb{N}^{2n}$, there exists $C_\alpha > 0$, such that

$$\|\nabla_{x,\xi}^\alpha a(x, \xi; h)\| \leq C_\alpha m(x, \xi) h^{-p}.$$

For a symbol $a(x, \xi; h)$, we define the Weyl quantization, $a^w(x, h \nabla_x; h) := Op_h^\omega(a)$ by

$$Op_h^\omega(a)u(x) = \frac{1}{(2\pi h)^n} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} e^{ih^{-1}(x-y) \cdot \xi} a\left(\frac{x+y}{2}, \xi; h\right) u(y) dy d\xi,$$

where $u(x)$ is in the Schwarz space.

5.1. Construction of $\widehat{H}_\theta$. We follow the approach of Sjöstrand [39]. Let Ω be a complex domain satisfying the assumption $(\mathbf{A}_\Omega^+)$ and $\psi \in C_0^\infty(\mathbb{R}^3)$ be such that $\psi(x) \geq 0$, $\psi(x) = 1$ if $|x| \leq 1$ and $\psi(x) = 0$ if $|x| \geq 2$. We recall the notations of Section 3: $\phi_\theta(x) = x + \theta g(x)$ with $g(x) = 0$ in the compact set $B(0, R_0) \subset K$ and $g(x) = x$ outside $K \subset B(0, \alpha_0)$ where $\alpha_0 > 0$ is sufficiently large.

Using Lemma 3, the semi-classical principal symbol of H_θ is given by:

$$h_\theta(x, \xi) = \alpha \cdot \zeta_\theta(x, \xi) + mc^2\beta + V(\phi_\theta(x)) - c\frac{1}{2} \sum_{j=1}^3 \alpha_j \cdot J_{\phi_\theta(x)}^{-1} \partial_j J_{\phi_\theta(x)},$$

with

$$\zeta_\theta(x, \xi) = (\zeta_{\theta,1}(x, \xi), \zeta_{\theta,2}(x, \xi), \zeta_{\theta,3}(x, \xi)) \text{ and } \zeta_{\theta,j}(x, \xi) = c \sum_{k=1}^3 \xi_k \left(\partial_j \phi_{\theta,k}^{-1} \right) (\phi_\theta(x)).$$

For all (x, ξ) , the matrix $M = \alpha \cdot \zeta_\theta(x, \xi) + mc^2\beta$, has two eigenvalues

$$\lambda_\theta^\pm = \pm \sqrt{\zeta_\theta(x, \xi)^2 + m^2 c^4}.$$

Consequently, there exists an invertible matrix U such that

$$U^{-1} M U = d_\theta := \begin{pmatrix} \lambda_\theta^+ I_2 & 0 \\ 0 & \lambda_\theta^- I_2 \end{pmatrix},$$

where

$$U = \begin{pmatrix} I_2 & \frac{-1}{\lambda_\theta^+ + mc^2} \sigma \cdot \zeta_\theta(x, \xi) \\ \frac{1}{\lambda_\theta^+ + mc^2} \sigma \cdot \zeta_\theta(x, \xi) & I_2 \end{pmatrix},$$

with $\sigma = (\sigma_1, \sigma_2, \sigma_3)$ and $(\sigma_j)_{1 \leq j \leq 3}$ the 2×2 Pauli matrices.

One can easily prove that the norms of U , U^{-1} and their derivatives are bounded in the following way:

$$(20) \quad \begin{aligned} \|\partial_x^\alpha \partial_\xi^\beta U(x, \xi)\| &< C \langle \xi \rangle^{-\beta}, \\ \|\partial_x^\alpha \partial_\xi^\beta U^{-1}(x, \xi)\| &< C \langle \xi \rangle^{-\beta}, \quad \forall \alpha, \beta \in \mathbb{N}. \end{aligned}$$

Applying U^{-1} (resp. U) on the left (resp. on the right) of $h_\theta(x, \xi)$, we obtain

$$U^{-1} h_\theta(x, \xi) U = d_\theta + \tilde{V}_\theta(x, \xi),$$

where $\tilde{V}_\theta(x, \xi) = U^{-1} \left(V(\phi_\theta(x)) - c\frac{1}{2} \sum_{j=1}^3 \alpha_j \cdot J_{\phi_\theta(x)}^{-1} \partial_j J_{\phi_\theta(x)} \right) U$. Since the term $\partial_j (J_{\phi_\theta(x)})$ is compactly supported $j=1, 2, 3$, (see Lemma 3) and the matrix $V(\phi_\theta(x))$, U , U^{-1} , and their derivatives are uniformly bounded, then $\tilde{V}_\theta(x, \xi)$ and their derivatives are uniformly bounded.

In order to construct $\widehat{H}_\theta$, we introduce an intermediate function $f(x, \xi)$:

We denote $|\Omega|$ the diameter of Ω . Let us choose $\beta_0 > 0$ and $C_0 > 0$, sufficiently large such that

$$(21) \quad \begin{aligned} \forall \xi \in \mathbb{R}^3; \quad \sup_{x \in \mathbb{R}^3} \|\tilde{V}_\theta(x, \xi)\| + |\Omega| &\leq \frac{1}{2} |\lambda_\theta^\pm - iC_0\psi(\frac{\xi}{\beta_0})| \\ &= \frac{1}{2} \sqrt{(\operatorname{Re}(\lambda_\theta^\pm))^2 + \left(\operatorname{Im}(\lambda_\theta^\pm) - C_0\psi(\frac{\xi}{\beta_0}) \right)^2}. \end{aligned}$$

We prove (21) considering the two cases:

- For $|\xi| > \beta_0$, with $\beta_0 > 0$, sufficiently large we have

$$\sup_{x \in \mathbb{R}^3} \|\tilde{V}_\theta(x, \xi)\| + |\Omega| \leq \frac{1}{2} |\operatorname{Re}(\lambda_\theta^\pm)|.$$

- For $|\xi| \leq \beta_0$, since $\lambda_\theta^\pm$ is bounded, we choose $C_0 > 0$, sufficiently large such that

$$\sup_{x \in \mathbb{R}^3} \|\tilde{V}_\theta(x, \xi)\| + |\Omega| \leq \frac{1}{2} |\operatorname{Im}(\lambda_\theta^\pm) - C_0|.$$

For $|x| > \alpha_0 > 0$, sufficiently large we have $\zeta_{\theta,j}(x, \xi) = \frac{c\xi_j}{1+\theta}$ and $\lambda_\theta^\pm = \pm c\sqrt{\frac{\xi^2}{(1+\theta)^2} + m^2c^2}$. Since the domain Ω satisfies the assumption $(\mathbf{A}_\Omega^+)$, we have

$$\min\{\operatorname{dist}(\overline{\Omega}, \lambda_\theta^+), \operatorname{dist}(\overline{\Omega}, \lambda_\theta^-)\} \neq 0,$$

hence we can choose $\alpha_0 > 0$ sufficiently large such that

$$(22) \quad \forall |x| > \alpha_0, \quad \|\tilde{V}_\theta(x, \xi)\| \leq \frac{1}{2} \operatorname{dist}(\overline{\Omega}, \lambda_\theta^\pm) := \frac{1}{2} \min\{\operatorname{dist}(\overline{\Omega}, \lambda_\theta^+), \operatorname{dist}(\overline{\Omega}, \lambda_\theta^-)\}.$$

Now, we define $f(x, \xi)$ in the following way:

$$(23) \quad f(x, \xi) = C_0 \psi\left(\frac{x}{\alpha_0}\right) \psi\left(\frac{\xi}{\beta_0}\right).$$

Lemma 6. *The matrix $h_\theta(x, \xi) - if(x, \xi) - z$ is invertible for all $z \in \Omega$ and satisfies*

$$(24) \quad \|\partial_x^\alpha \partial_\xi^\beta (h_\theta(x, \xi) - if(x, \xi) - z)^{-1}\| < C\langle \xi \rangle^{-1-\beta}, \quad \forall \alpha, \beta \in \mathbb{N}.$$

Proof. Applying U^{-1} (resp. U) on the left (resp. on the right) of $h_\theta(x, \xi) - if(x, \xi) - z$, we obtain

$$U^{-1} (h_\theta(x, \xi) - if(x, \xi) - z) U = d_\theta - if(x, \xi) - z + \tilde{V}_\theta(x, \xi).$$

1) Let us prove that the symbol $\sigma := d_\theta - if(x, \xi) - z + \tilde{V}_\theta(x, \xi)$ is invertible.

- For $|x| \leq \alpha_0$,

$$\sigma = (d_\theta - iC_0\psi(\frac{\xi}{\beta_0})) \left(I_4 + \left(d_\theta - iC_0\psi(\frac{\xi}{\beta_0}) \right)^{-1} (\tilde{V}_\theta(x, \xi) - z) \right).$$

According to (21), we have

$$\left\| \left(d_\theta - iC_0\psi(\frac{\xi}{\beta_0}) \right)^{-1} (\tilde{V}_\theta(x, \xi) - z) \right\| < \frac{1}{2},$$

thus σ is invertible and satisfies

$$(25) \quad \|\sigma^{-1}\| < 2 \left\| \left(d_\theta - iC_0\psi(\frac{\xi}{\beta_0}) \right)^{-1} \right\| < C\langle \xi \rangle^{-1}.$$

- For $|x| > \alpha_0$, we have $\lambda_\theta^\pm = \pm c\sqrt{\frac{\xi^2}{(1+\theta)^2} + m^2c^2}$. Since $f(x, \xi) \geq 0$, we have

$$|\lambda_\theta^+ - (z + if(x, \xi))| > \operatorname{dist}(\overline{\Omega}, \lambda_\theta^\pm) > C\langle \xi \rangle > 0,$$

and

$$|\operatorname{Re}(\lambda_\theta^- - (z + if(x, \xi)))| = |\operatorname{Re}(\lambda_\theta^- - z)| > \operatorname{dist}(\overline{\Omega}, \lambda_\theta^\pm) > C\langle \xi \rangle > 0.$$

Since

$$\sigma = (d_\theta - if(x, \xi) - z) \left(I_4 + (d_\theta - if(x, \xi) - z)^{-1} \tilde{V}_\theta(x, \xi) \right),$$

and

$$\| (d_\theta - if(x, \xi) - z)^{-1} \tilde{V}_\theta(x, \xi) \| < \frac{1}{2}, \quad (\text{see (22)}),$$

the matrix σ is invertible and

$$(26) \quad \|\sigma^{-1}\| < 2\| (d_\theta - if(x, \xi) - z)^{-1} \| < C\langle \xi \rangle^{-1}.$$

2) According to 1), the matrix $U^{-1}(h_\theta(x, \xi) - if(x, \xi) - z)U$ is invertible. From (25), (26) and (20), we deduce that the matrix $h_\theta(x, \xi) - if(x, \xi) - z$ is invertible and

$$(27) \quad \begin{aligned} \| (h_\theta(x, \xi) - if(x, \xi) - z)^{-1} \| &= \| U^{-1} \left(d_\theta - if(x, \xi) I_4 - z + \tilde{V}_\theta(x, \xi) \right)^{-1} U \| \\ &\leq \| U \| \| U^{-1} \| \| \left(d_\theta - if(x, \xi) I_4 - z + \tilde{V}_\theta(x, \xi) \right)^{-1} \| \\ &< C\langle \xi \rangle^{-1}. \end{aligned}$$

This gives (24) for $\alpha = \beta = 0$. Using (20) and (27) we obtain (24) for $(\alpha, \beta) \in \mathbb{N}^2$ by induction. $\square$

We denote $\tilde{H}_\theta = H_\theta + \tilde{T}$, with $\tilde{T} = Op_h^\omega(-if(x, \xi))$, where $f(x, \xi)$ is defined in (23). It's clear that the semi-classical principal symbol of $(\tilde{H}_\theta - z)$ is

$$\sigma_{\tilde{H}_\theta} := h_\theta(x, \xi) - if(x, \xi) - z.$$

Proposition 5. *If $h > 0$ is small enough, the operator $(z - \tilde{H}_\theta)$ is invertible for every $z \in \Omega$ and, for every $N \in \mathbb{N}$ its inverse satisfies:*

$$(z - \tilde{H}_\theta)^{-1} = O_N(1) : D(H^N) \mapsto D(H^{N+1}),$$

uniformly for $z \in \Omega$. Here $D(H^N)$ designates the domain of H^N with the convention $D(H^0) = \mathcal{H}$.

Proof. Let us prove that the operator $(z - \tilde{H}_\theta)$ is a Fredholm operator of index 0. We have

$$\begin{aligned} (z - \tilde{H}_\theta)(z - H_{0,\theta})^{-1} &= (z - H_{0,\theta} + H_{0,\theta} - \tilde{H}_\theta)(z - H_{0,\theta})^{-1} \\ &= I - (\tilde{T} + V(\phi_\theta(x)))(z - H_{0,\theta})^{-1}. \end{aligned}$$

Since the right-hand side is a perturbation of the identity by a compact operator and

$$(z - H_{0,\theta})^{-1} : (L^2(\mathbb{R}^3))^4 \mapsto D(H) \text{ is invertible,}$$

the operator $(z - \tilde{H}_\theta)$ is Fredholm of index 0. Consequently, it is enough to show that

$$(28) \quad \|u\|_{D(H^{N+1})}^2 \leq C\|(z - \tilde{H}_\theta)u\|_{D(H^N)}^2, \quad \text{for } u \in D(H^{N+1}).$$

According to Lemma 6, the symbol $q_0 = \sigma_{\tilde{H}_\theta}^{-1}$ is well defined and satisfies

$$\|\partial_x^\alpha \partial_\xi^\beta q_0\| < C\langle \xi \rangle^{-1-\beta}.$$

Moreover, having

$$\|\partial_x^\alpha \partial_\xi^\beta (\sigma_{\tilde{H}_\theta})\| < C\langle \xi \rangle^{+1-\beta},$$

the composition theorem of h -pseudo-differential operators implies

$$Op_h^\omega(q_0)Op_h^\omega(\sigma_{\tilde{H}_\theta}) = Op_h^\omega(r)$$

where $(r - 1)$ is in the space of symbols $\mathcal{S}^0(h)$. In particular the operator

$$Op_h^\omega(r) : D(H^{N+1}) \mapsto D(H^{N+1}), \quad \forall N \in \mathbb{N},$$

is invertible for h small enough, then (28) follows. Therefore the operator $(\tilde{H}_\theta - z)$ is also invertible and we have

$$(z - \tilde{H}_\theta)^{-1} = O_N(1) : D(H^N) \mapsto D(H^{N+1}).$$

□

Proposition 6. *There exists $\hat{H}_\theta : D(H) \mapsto \mathcal{H}$, with the following properties.*

The difference $K := \hat{H}_\theta - H_\theta$ is of finite rank $O(h^{-3})$, has a compact support in the sense that $K = \chi_1 K \chi_1$ for some $\chi_1 \in C_0^\infty(\mathbb{R}^3)$ and

$$K = O(1) : D(H^N) \mapsto D(H^M) \quad \forall N, M \in \mathbb{N}.$$

Moreover, for every $N \in \mathbb{N}$, we have

$$(\hat{H}_\theta - z)^{-1} = O(1) : D(H^N) \mapsto D(H^{N+1}),$$

uniformly for $z \in \overline{\Omega}$.

Proof. (We again use all the previous notations) We define

$$\hat{H}_\theta := H_\theta + \chi_1 T \chi_1 = \tilde{H}_\theta + \chi_1 T \chi_1 - \tilde{T},$$

with $\chi_1(x) = \psi(\frac{x}{2\alpha_0})$ and

$$T := \chi(-h^2\Delta + x^2)\tilde{T} = \chi(-h^2\Delta + x^2)Op_h^\omega(-if(x, \xi))$$

where $\chi \in C_0^\infty(\mathbb{R})$ is such that:

$$\chi(\xi^2 + x^2) = 1 \text{ on the support of } f(x, \xi) \text{ (see (23)).}$$

By functional calculus (see [12]), we can prove that

$$(29) \quad \tilde{H}_\theta - \hat{H}_\theta = \tilde{T} - \chi_1 T \chi_1 = O(h^\infty) : D(H^N) \mapsto D(H^M), \quad \forall M, N \in \mathbb{N}.$$

The last lemma, the formula (29) and

$$(\hat{H}_\theta - z)^{-1} = (\tilde{H}_\theta - z)^{-1} \left(I + (\hat{H}_\theta - \tilde{H}_\theta)(\tilde{H}_\theta - z)^{-1} \right)^{-1}$$

yield for all $N \in \mathbb{N}$

$$(\hat{H}_\theta - z)^{-1} = O(1) : D(H^N) \mapsto D(H^{N+1}).$$

According to the fact that $\chi(-h^2\Delta + x^2)$ is of finite rank $O(h^{-3})$, to the fact that Weyl quantification $Op_h^\omega(-if(x, \xi))$ is bounded and to the definition of χ_1 , the operator

$$K := \hat{H}_\theta - H_\theta = \chi_1 (\chi(-h^2\Delta + x^2)Op_h^\omega(-if(x, \xi))) \chi_1$$

is of finite rank $O(h^{-3})$ and compactly supported. □

5.2. Upper bound for the number of resonances. In this section we establish the upper bound of the number of resonances given in Theorem 1.

Lemma 7. *Let $\rho > 0$, Ω be an open complex relatively compact subset of $\mathbb{C}$ and H_θ be define as above. There exists g satifying $(\mathbf{A}_g)$ such that for h small enough and $z \in \overline{\Omega} \cap \{\text{Im } z \geq \rho > 0\}$, we have $(z - H_\theta)^{-1} = O(1)$.*

Proof. We again use the notations of Section 3: $\phi_\theta(x) = x + \theta g(x)$ with $g(x) = 0$ in the compact $B(0, R_0)$, and the notations of Subsection 5.1 concerning $h_\theta(x, \xi)$, U , U^{-1} , d_θ and $\tilde{V}_\theta(x, \xi)$ which satisfy

$$U^{-1}h_\theta(x, \xi)U = d_\theta + \tilde{V}_\theta(x, \xi).$$

The matrice $h_\theta(x, \xi)$ is the semi-classical principal symbol of H_θ .

According to Section 4, the resonances are independent of the family U_θ . Then we can assume that $g(x) = 0$ in the ball $B(0, R_g) \supset B(0, R_0)$, with $R_g > 0$, sufficiently large such that

$$\forall x \in \mathbb{R}^3, \quad |x| > R_g > 0, \quad \|\tilde{V}_\theta(x, \xi)\| \leq \frac{\rho}{2}.$$

Repeating arguments of Subsection 5.1, we can prove that $(d_\theta + \tilde{V}_\theta(x, \xi) - z)$ is invertible, then $(h_\theta(x, \xi) - z)$ is invertible and

$$\|\partial_x^\alpha \partial_\xi^\beta (h_\theta(x, \xi) - z)^{-1}\| < C \langle \xi \rangle^{-1-\beta}.$$

Since we have:

$$\|\partial_x^\alpha \partial_\xi^\beta (h_\theta(x, \xi) - z)\| < C \langle \xi \rangle^{+1-\beta},$$

the composition theorem of h -pseudo-differential operators implies

$$Op_h^\omega((h_\theta(x, \xi) - z)^{-1})Op_h^\omega(h_\theta(x, \xi) - z) = 1 + O(h),$$

where $O(h)$ corresponds to the norm in $\mathcal{L}(L^2)$. □

Proof of the Theorem 1.

Let $\hat{K}(z) = K(z - \hat{H}_\theta)^{-1}$ with $\hat{H}_\theta$, K defined in Proposition 6. We remark that

$$(I + \hat{K}(z))(z - \hat{H}_\theta) = (z - \hat{H}_\theta) + K = z - H_\theta.$$

Thus, the resonances $z \in \text{Res}(H) \cap \Omega$ repeated with their multiplicities coincide with the zeros of the function

$$D(z) = \det(I + \hat{K}(z)).$$

Since K is bounded and is of finite rank $O(h^{-3})$, we have

$$|D(z)| \leq e^{\|\hat{K}(z)\|_{tr}} \leq e^{C_0 h^{-3}}, \quad \text{for all } z \in \overline{\Omega}.$$

Using Lemma 7, we get $(z - H_\theta)^{-1} = O(1)$ for $\text{Im } z \geq \rho > 0$ and $z \in \overline{\Omega}$. Since

$$(30) \quad (I + \hat{K}(z))^{-1} = (z - \hat{H}_\theta)(z - H_\theta)^{-1},$$

then

$$\|(I + \hat{K}(z))^{-1}\| \leq C_1, \quad \text{Im } z \geq \rho > 0.$$

Writing the operator $(I + \hat{K}(z))^{-1}$ in the form

$$(I + \hat{K}(z))^{-1} = I - \hat{K}(z)(I + \hat{K}(z))^{-1},$$

we obtain

$$|\det \left((I + \widehat{K}(z))^{-1} \right)| \leq e^{C_2 h^{-3}}, \quad \text{Im } z \geq \rho,$$

which implies

$$|D(z)| \geq C e^{-C_3 h^{-3}}, \quad z \in \overline{\Omega} \cap \{\text{Im } z \geq \rho\}.$$

Now, applying Jensen's inequality in a slightly larger domain, we obtain Theorem 1. $\square$

6. REPRESENTATION OF DERIVATIVE OF THE SPECTRAL SHIFT FUNCTION

In this section we prove our principal result given in Theorem 2 and a generalization (see Theorem 5). Moreover, we give a Sjöstrand type local trace formula.

The spectral shift function $\xi(\lambda, h) \in \mathcal{D}'(\mathbb{R})$ associated to H_0, H_1 is defined (see [10], [36], [44]) by

$$\langle \xi'(\lambda, h), f(\lambda) \rangle = \text{tr}(f(H_1) - f(H_0)), \quad f \in C_0^\infty(\mathbb{R}).$$

In the following, we will use the notations:

$$H_1 = H, \quad K_1 := K = \widehat{H}_{1,\theta} - H_{1,\theta} := \widehat{H}_\theta - H_\theta \text{ and } [a.]_0^1 = a_1 - a_0.$$

For an integer $m > 3$, we define the functions:

$$(31) \quad \sigma_\pm(z) = (z^2 + 1)^m \text{tr} \left[(H. - i)^{-m} (H. + i)^{-m} (z - H.)^{-1} \right]_0^1, \quad \pm \text{Im } z > 0.$$

The $\sigma_\pm$ satisfy the relation

$$(32) \quad \sigma_-(z) = \overline{\sigma_+(\bar{z})}, \quad \text{Im}(z) < 0.$$

Proposition 7. *For a potential V satisfying the assumption $(\mathbf{A}_V)$ with $\delta > 3$, the function $\theta \mapsto \left[(H_{.,\theta} - i)^{-m} (H_{.,\theta} + i)^{-m} (z - H_{.,\theta})^{-1} \right]_0^1$ is holomorphic from D_ϵ^+ to the space of trace class operators. Moreover, for any $\theta \in D_\epsilon^+$, we have*

$$(33) \quad \sigma_\pm(z) = (z^2 + 1)^m \text{tr} \left[(H_{.,\theta} - i)^{-m} (H_{.,\theta} + i)^{-m} (z - H_{.,\theta})^{-1} \right]_0^1, \quad \pm \text{Im } z > 0.$$

Proof. For $\theta \in \mathbb{R}$, the operator

$$(H. - i)^{-m} (H. + i)^{-m} (z - H.)^{-1},$$

is unitarily equivalent to the operator

$$(H_{.,\theta} - i)^{-m} (H_{.,\theta} + i)^{-m} (z - H_{.,\theta})^{-1}.$$

Using the cyclicity of the trace, we deduce

$$(34) \quad \sigma_\pm(z) = (z^2 + 1)^m \text{tr} \left[(H_{.,\theta} - i)^{-m} (H_{.,\theta} + i)^{-m} (z - H_{.,\theta})^{-1} \right]_0^1, \quad \pm \text{Im } z > 0, \quad \theta \in \mathbb{R}.$$

According to the proof of Theorem 4, the resolvent $(z - H_{.,\theta})^{-1}$ is analytic for $\theta \in D_\epsilon^+$ and $z \in \Omega \cap \{\text{Im } z > 0\}$. Then, the function $\theta \mapsto (H_{.,\theta} - i)^{-m} (H_{.,\theta} + i)^{-m} (z - H_{.,\theta})^{-1}$ is also analytic on D_ϵ^+ .

Now, we treat the difference

$$(35) \quad \begin{aligned} \left[(z - H_{.,\theta})^{-1} (H_{.,\theta} - i)^{-m} (H_{.,\theta} + i)^{-m} \right]_0^1 &= A_1 B_1 C_1 - A_0 B_0 C_0 \\ &= A_1 B_1 (C_1 - C_0) + A_1 (B_1 - B_0) C_0 \\ &\quad + (A_1 - A_0) B_0 C_0. \end{aligned}$$

Clearly, the terms $A := (z - H_{\cdot,\theta})^{-1}$ for $\text{Im } z > 0$, $B := (H_{\cdot,\theta} - i)^{-m}$ and $C := (H_{\cdot,\theta} + i)^{-m}$ are bounded.

For any integer $m > 3$, the term

$$(36) \quad B_1(C_1 - C_0) = \left(B_1(C_1 - C_0) \langle x \rangle^\delta \langle h \nabla_x \rangle^m \right) \left(\langle h \nabla_x \rangle^{-m} \langle x \rangle^{-\delta} \right),$$

is analytic for $\theta \in D_\epsilon^+$ with values in the space of trace class operators. This can be proved using functional calculus in the framework of h -pseudo-differential operators (see [12]): The first factor $B_1(C_1 - C_0) \langle x \rangle^\delta \langle h \nabla_x \rangle^m$ is analytic for $\theta \in D_\epsilon^+$, the second factor $(\langle h \nabla_x \rangle^{-m} \langle x \rangle^{-\delta})$ is in the space of trace class operators and its trace norm is bounded by $O(h^{-3})$. Then, the left-hand side of the equation (36) is in the space of trace class operators and its trace norm is bounded by $O(h^{-3})$. The same argument can be used for the terms $A_1(B_1 - B_0)$ and $(A_1 - A_0)B_0$, then their trace norm are bounded by $O(h^{-3})$.

Since the function $\text{tr} \left[(H_{\cdot,\theta} - i)^{-m} (H_{\cdot,\theta} + i)^{-m} (z - H_{\cdot,\theta})^{-1} \right]_0^1$ is analytic with respect to $\theta \in D_\epsilon^+$ and independent of θ on the real axis, the formula (33) follows. $\square$

Repeating the construction of $\widehat{H}_{1,\theta}$, we can construct an operator $\widehat{H}_{0,\theta} : D(H_0) \rightarrow \mathcal{H}$ with the properties of $\widehat{H}_{0,\theta}$ such that the difference $K_0 := \widehat{H}_{0,\theta} - H_{0,\theta}$ satisfies the properties of K_1 (see Proposition 6).

Proposition 8. *There exists a function $a_+(z, h)$ holomorphic in Ω , such that for all $z \in \Omega \cap \{\text{Im}(z) > 0\}$, we have:*

$$(37) \quad \sigma_+(z) = \text{tr} \left[(H_{\cdot,\theta} - z)^{-1} K. (\widehat{H}_{\cdot,\theta} - z)^{-1} \right]_0^1 + a_+(z, h),$$

$$|a_+(z, h)| \leq C(\Omega) h^{-3}, \quad z \in \Omega,$$

with $C(\Omega)$ a constant independent of h .

Proof. For $z \in \Omega \cap \{\text{Im} z > 0\}$, we have

$$(38) \quad (H_{\cdot,\theta} - z)^{-1} = (\widehat{H}_{\cdot,\theta} - z)^{-1} + (H_{\cdot,\theta} - z)^{-1} K. (\widehat{H}_{\cdot,\theta} - z)^{-1}.$$

From the equations (38) and (33), we deduce:

$$\begin{aligned} \sigma_+(z) &= ((z - i)(z + i))^m \text{tr} \left[\left((\widehat{H}_{\cdot,\theta} - z)^{-1} (H_{\cdot,\theta} - i)^{-m} (H_{\cdot,\theta} + i)^{-m} \right) \right]_0^1 \\ &+ ((z - i)(z + i))^m \text{tr} \left[\left((H_{\cdot,\theta} - z)^{-1} K. (\widehat{H}_{\cdot,\theta} - z)^{-1} (H_{\cdot,\theta} - i)^{-m} (H_{\cdot,\theta} + i)^{-m} \right) \right]_0^1 \\ &= A(z) + B(z). \end{aligned}$$

Starting with the resolvent equation, we obtain:

$$\begin{aligned} &((z - i)(z + i))^m (H_{\cdot,\theta} - i)^{-m} (H_{\cdot,\theta} + i)^{-m} (H_{\cdot,\theta} - z)^{-1} \\ &= (H_{\cdot,\theta} - z)^{-1} - \sum_{k=1}^m (z + i)^{k-1} (H_{\cdot,\theta} + i)^{-k} \\ &- (z + i) \sum_{k=1}^m (z - i)^{k-1} (H_{\cdot,\theta} + i)^{-m} (H_{\cdot,\theta} - i)^{-k}. \end{aligned}$$

Using the last equation, the cyclicity of the trace and Proposition 6 we obtain

$$\begin{aligned} B(z) &= \operatorname{tr} \left[K.(\widehat{H}_{\cdot,\theta} - z)^{-1} \left((H_{\cdot,\theta} - z)^{-1} - \sum_{k=1}^m (z+i)^{k-1} (H_{\cdot,\theta} + i)^{-k} \right. \right. \\ &\quad \left. \left. - (z+i) \sum_{k=1}^m (z-i)^{k-1} (H_{\cdot,\theta} + i)^{-m} (H_{\cdot,\theta} - i)^{-k} \right) \right]_0^1 \\ &= \operatorname{tr} \left[(H_{\cdot,\theta} - z)^{-1} K.(\widehat{H}_{\cdot,\theta} - z)^{-1} \right]_0^1 + b(z). \end{aligned}$$

Since the operator $(\widehat{H}_{\cdot,\theta} - z)^{-1}$ is bounded and holomorphic in Ω by construction, $b(z)$ is holomorphic and bounded by $O(h^{-3})$.

It remains to show that

$$\begin{aligned} A(z) &= ((z-i)(z+i))^m \operatorname{tr} \left[(\widehat{H}_{\cdot,\theta} - z)^{-1} (H_{\cdot,\theta} - i)^{-m} (H_{\cdot,\theta} + i)^{-m} \right]_0^1 \\ &= ((z-i)(z+i))^m \operatorname{tr} (\widehat{A}_1 B_1 C_1 - \widehat{A}_0 B_0 C_0), \end{aligned}$$

is holomorphic and bounded by $O(h^{-3})$.

We recall that the terms $\widehat{A}_{\cdot} := (\widehat{H}_{\cdot,\theta} - z)^{-1}$ for $z \in \Omega$, $B_{\cdot} := (H_{\cdot,\theta} - i)^{-m}$ and $C_{\cdot} := (H_{\cdot,\theta} + i)^{-m}$ are bounded. Using the assumption $(\mathbf{A}_{\mathbf{V}})$ with $\delta > 3$, we treat the difference $(\widehat{A}_1 B_1 C_1 - \widehat{A}_0 B_0 C_0)$ as (35). The only difference is for the term $(\widehat{A}_1 - \widehat{A}_0) B_0$. We write

$$\begin{aligned} (\widehat{A}_1 - \widehat{A}_0) B_0 &= (\widehat{H}_{1,\theta} - z)^{-1} (\widehat{H}_{0,\theta} - \widehat{H}_{1,\theta}) (\widehat{H}_{0,\theta} - z)^{-1} (H_{0,\theta} - i)^{-m}, \\ \text{with } \widehat{H}_{0,\theta} - \widehat{H}_{1,\theta} &= H_{0,\theta} - H_{1,\theta} + K_0 - K_1. \end{aligned}$$

Then, modulo a trace class operator uniformly bounded, with trace norm bounded by $O(h^{-3})$, we have

$$\begin{aligned} (\widehat{A}_1 - \widehat{A}_0) B_0 &= (\widehat{H}_{1,\theta} - z)^{-1} \circ ((H_{0,\theta} - H_{1,\theta})(H_{1,\theta} - i)^{-m}) \\ &\quad \circ \left((H_{1,\theta} - i)^m (\widehat{H}_{0,\theta} - z)^{-1} (H_{0,\theta} - i)^{-m} \right). \end{aligned}$$

The second factor $(H_{0,\theta} - H_{1,\theta})(H_{1,\theta} - i)^{-m}$ is trace class and its trace is $O(h^{-3})$, the first and the third factor are bounded. Then, the term $(\widehat{A}_1 - \widehat{A}_0) B_0$ is analytic for $z \in \Omega$ with values in the space of trace class operators and its trace is bounded by $O(h^{-3})$ and so is the difference $(\widehat{A}_1 B_1 C_1 - \widehat{A}_0 B_0 C_0)$. $\square$

Lemma 8. *For $f \in C_0^\infty(\mathbb{R})$, we have*

$$(39) \quad \langle \xi', f \rangle = \lim_{\varepsilon \rightarrow 0} \frac{i}{2\pi} \int f(\lambda) [\sigma_+(\lambda + i\varepsilon) - \sigma_-(\lambda - i\varepsilon)] d\lambda.$$

This limit is taken in the sense of distribution.

Proof. We follow the proof of [13, Lemma 1]. Let $f \in C_0^\infty(\mathbb{R})$, $\tilde{f}(z) \in C_0^\infty(\mathbb{R}^2)$ be an almost analytic extension of f and

$$g(x) = f(x)(x^2 + 1)^m.$$

Then

$$g(H_{\cdot}) = -\frac{1}{\pi} \int \bar{\partial}_z \tilde{f}(z) (z^2 + 1)^m (z - H_{\cdot})^{-1} L(dz),$$

where $L(dz)$ is the Lebesgue measure on $\mathbb{C}$. Clearly

$$\begin{aligned} f(H.) &= (H. - i)^{-m}(H. + i)^{-m}g(H.) \\ &= -\frac{1}{\pi} \int \bar{\partial}_z \tilde{f}(z)(z^2 + 1)^m (H. - i)^{-m}(H. + i)^{-m}(z - H.)^{-1} L(dz) \end{aligned}$$

which implies:

$$\begin{aligned} \text{tr}(f(H_1) - f(H_0)) &= -\frac{1}{\pi} \int \bar{\partial}_z \tilde{f}(z)(z^2 + 1)^m \\ (40) \quad &\times \text{tr} \left[(H. - i)^{-m}(H. + i)^{-m}(z - H.)^{-1} \right]_0^1 L(dz). \end{aligned}$$

We have $\sigma_{\pm}(z) = O(h^{-3}|\text{Im } z|^{-2})$ and the derivative $\bar{\partial}_z \tilde{f} = O(|\text{Im } z|^N)$ for all $N \in \mathbb{N}$ ($f \in C_0^\infty(\mathbb{R})$), so we write the right-hand side of (40) as

$$\begin{aligned} \langle \xi', f \rangle &= \text{tr}(f(H_1) - f(H_0)) \\ &= -\frac{1}{\pi} \lim_{\varepsilon \rightarrow 0} \left(\int_{\text{Im } z > 0} \bar{\partial}_z \tilde{f}(z) \sigma_+(z + i\varepsilon) L(dz) + \int_{\text{Im } z < 0} \bar{\partial}_z \tilde{f}(z) \sigma_-(z - i\varepsilon) L(dz) \right). \end{aligned}$$

According to Proposition 8, the functions $\sigma_+(z + i\varepsilon)$ and $\sigma_-(z - i\varepsilon)$ are holomorphic in $\{z \in \Omega; \text{Im } z > 0\}$ and $\{z \in \Omega; \text{Im } z < 0\}$ respectively. Applying the Green formula, we obtain the lemma. $\square$

Before the proof of Theorem 2, let us give the following proposition:

Proposition 9. (see [38], [39]) *Let $F(z, h)$ be a holomorphic function in an open simply connected domain Ω containing a number $N(h)$ of zeros. We suppose that,*

$$F(z, h) = O(1)e^{O(1)N(h)}, \quad z \in \Omega,$$

and for all $\rho > 0$ small enough, there exists $C > 0$ such that for all $z \in \Omega_\rho := \Omega \cap \{\text{Im } z > \rho\}$ we have

$$|F(z, h)| \geq e^{-CN(h)}.$$

Then for all open simply connected subset $\tilde{\Omega} \Subset \Omega$ there exists $g(., h)$ holomorphic in $\tilde{\Omega}$ such that

$$F(z, h) = \prod_{j=1}^{N(h)} (z - z_j) e^{g(z, h)}, \quad \partial_z g(z, h) = O(N(h)), \quad z \in \tilde{\Omega}.$$

Proof of Theorem 2. We follow the argument of Sjöstrand ([39]). Let

$$\hat{K}_.(z) = K_.(z - \hat{H}_.)_{,\theta}^{-1}.$$

As [39, Equation(4.31)] we have the representation

$$\begin{aligned} -\text{tr} \left((H_.)_{,\theta} - z \right)^{-1} K_.(\hat{H}_.)_{,\theta}^{-1} &= \text{tr} \left(\left(1 + \hat{K}_.(z) \right)^{-1} \partial_z \hat{K}_.(z) \right) \\ &= \partial_z \log \det \left(1 + \hat{K}_.(z) \right). \end{aligned}$$

From Subsection 5.2 the resonances are the zeros of the function

$$D(z, h) = \det \left(I + \hat{K}_1(z) \right) = O(1)e^{ch^{-3}}.$$

Since the function $\det(1 + \widehat{K}_0(z))$ has no zeros in Ω (see (30) and Remark 6), the term $\partial_z \log \det(1 + \widehat{K}_0(z))$ is analytic and using the Proposition 9, it is bounded by $O(h^{-3})$.

We recall the $\text{Res}(H)$ be the set of resonances of H and let

$$D(z, h) = G(z, h) \prod_{w \in \text{Res}(H) \cap \Omega} (z - w),$$

where, $G(z, h)$ and its inverse are holomorphic functions in Ω . Obviously,

$$(41) \quad \partial_z \log D(z, h) = \partial_z \log G(z, h) + \sum_{w \in \text{Res}(H) \cap \Omega} \frac{1}{z - w}.$$

Using Proposition 9, we have

$$|\partial_z \log G(z, h)| \leq C(\tilde{\Omega})h^{-3}, \quad z \in \tilde{\Omega},$$

where $\tilde{\Omega} \subset \subset \Omega$ is an open simply connected set and $C(\tilde{\Omega})$ is independent of h .

Now, we treat the non-holomorphic term in $(\sigma_+(\lambda + i\varepsilon) - \sigma_-(\lambda - i\varepsilon))$ when $\varepsilon \rightarrow 0$, which is

$$\sum_{w \in \text{Res}(H) \cap \Omega} \left(\frac{1}{\lambda + i\varepsilon - w} - \frac{1}{\lambda - i\varepsilon - \bar{w}} \right), \quad \text{for } \lambda \in I.$$

If $\text{Im}(w) \neq 0$, we have

$$\frac{-1}{2i\pi} \lim_{\varepsilon \rightarrow 0} \left(\frac{1}{\lambda + i\varepsilon - w} - \frac{1}{\lambda - i\varepsilon - \bar{w}} \right) = \frac{-\text{Im}(w)}{\pi|\lambda - w|^2},$$

while for $w \in \mathbb{R}$ we get

$$\frac{-1}{2i\pi} \lim_{\varepsilon \rightarrow 0} \left(\frac{1}{\lambda + i\varepsilon - w} - \frac{1}{\lambda - i\varepsilon - w} \right) = \delta(\lambda - w) = \delta_w(\lambda).$$

The second limit is taken in the sense of distributions.

Lemma 8 and Proposition 8 show that the function $r(z, h) = g(z, h) - \overline{g}(\bar{z}, h)$, with $g(z, h) = a_+(z, h) + \partial_z \log G(z, h) + \partial_z \log \det(1 + \widehat{K}_0(z))$ is a holomorphic function in Ω and satisfies the following estimate:

$$(42) \quad |g(z, h)| \leq C(\Omega)h^{-3}, \quad z \in W,$$

with $C(\Omega) > 0$ independent of h . □

Theorem 2 can be extended to a more general situation:

Theorem 5. *Assume that $H_1 = H_0 + V_1$, $H_2 = H_0 + V_2$. The potentials V_1, V_2 (resp. $V = V_1 - V_2$) satisfy the assumption $(\mathbf{A}_V)$ with $\delta > 0$ (resp. $\delta > 3$). Let Ω be a complex domain satisfying the assumption $(\mathbf{A}_\Omega^\pm)$, $W \Subset \Omega$ be an open simply connected and relatively compact set which is symmetric with respect to $\mathbb{R}$. Assume that $I = W \cap \mathbb{R}$ is an interval. Then for all $\lambda \in I$ we have a representation of the derivative of the spectral shift function*

associated to the pair (H_2, H_1) of the form:

$$(43) \quad \xi'(\lambda, h) = \frac{1}{\pi} \operatorname{Im} r(\lambda, h) + \left[\sum_{\substack{w \in \operatorname{Res}(H.) \cap \Omega \\ \operatorname{Im} w \neq 0}} \frac{-\operatorname{Im} w}{\pi |\lambda - w|^2} + \sum_{w \in \operatorname{Res}(H.) \cap I} \delta_w(\lambda) \right]_2^1,$$

where $r(z, h) = g(z, h) - \bar{g}(\bar{z}, h)$, $g(z, h)$ is a holomorphic function in Ω which satisfies the following estimate:

$$(44) \quad |g(z, h)| \leq C(W)h^{-3}, \quad z \in W,$$

with $C(W) > 0$ independent of h . Here $\delta_w(\cdot)$ is the Dirac mass at $w \in \mathbb{R}$.

Proof. We denote $H_{2,\theta} = U_\theta H_2 U_\theta^{-1}$ (U_θ defined in Section 3). As in Subsection 5.1, one constructs $\widehat{H}_{2,\theta} : D(H) \mapsto \mathcal{H}$ with the following properties:

$K_2 := \widehat{H}_{2,\theta} - H_{2,\theta}$ is of finite rank $O(h^{-3})$, has a compact support in the sense that $K_2 = \chi_2 K_2 \chi_2$ if $\chi_2 \in \mathbb{C}_0^\infty$ is equal to 1 on $B(0, R)$ for some sufficiently large R , and

$$(\widehat{H}_{2,\theta} - z)^{-1} = O(1) : \mathcal{H} \mapsto D(H), \text{ uniformly for } z \in \overline{\Omega}.$$

We repeat the proof of the Theorem 2 replacing K_0 by K_2 and $\widehat{K}_0(z)$ by $\widehat{K}_2(z) = K_2(z - \widehat{H}_{2,\theta})^{-1}$. Consequently $\partial_z \log \det (1 + \widehat{K}_0(z))$ is replaced by $\partial_z \log \det (1 + \widehat{K}_2(z))$ which is a non-holomorphic function. We treat this term as the term $\partial_z \log \det (1 + \widehat{K}_1(z))$ in the proof of Theorem 2. $\square$

Remark 8. The equation (6) shows that the spectral shift function $\xi(\lambda, h)$ satisfies

$$(45) \quad \begin{aligned} \xi(\lambda, h) - \xi(\lambda_0, h) &= \sum_{\substack{w \in \operatorname{Res}(H_1) \cap \Omega \\ \operatorname{Im} w \neq 0}} \frac{1}{\pi} \int_{\lambda_0}^{\lambda} \frac{-\operatorname{Im} w}{|\mu - w|^2} d\mu + \frac{1}{\pi} \int_{\lambda_0}^{\lambda} \operatorname{Im} r(\mu, h) d\mu \\ &+ \# \{ \mu \in [\lambda_0, \lambda]; \mu \in \sigma_d(H_1) \}. \end{aligned}$$

In particular, for $\lambda \in I \setminus \sigma_d(H_1)$ the distribution $\xi(\lambda, h)$ is continuous, and the function

$$\eta(\lambda, h) - \eta(\lambda_0, h) = \xi(\lambda, h) - \xi(\lambda_0, h) - \# \{ \mu \in [\lambda_0, \lambda]; \mu \in \sigma_d(H_1) \}$$

is real analytic in I . $\square$

Repeating the argument used in the proof of [8, Theorem 4], the following theorem is a direct consequence of Theorem 2.

Theorem 6. (Local trace formula) Let Ω be an open complex, simply connected and relatively compact set satisfying the assumption $(\mathbf{A}_\Omega^\pm)$ such that $I = \Omega \cap \mathbb{R}$ is an interval.

We suppose that f is a holomorphic function in $\overline{\Omega}$ and $\psi \in C_0^\infty(\mathbb{R})$ satisfies

$$\psi(\lambda) = \begin{cases} 0, & d(I, \lambda) > 2\varepsilon, \\ 1, & d(I, \lambda) < \varepsilon, \end{cases}$$

where $\varepsilon > 0$ and sufficiently small. Then

$$\operatorname{tr}[(\psi f)(H.)]_0^1 = \sum_{z \in \operatorname{Res}(H_1) \cap \Omega} f(z) + E_{\Omega, f, \psi}(h), \text{ with}$$

$$|E_{\Omega, f, \psi}(h)| \leq M(\psi, \Omega) \sup\{|f(z)|; 0 \leq d(\partial\Omega, z) \leq 2\varepsilon, \operatorname{Im}(z) \leq 0\} h^{-3}.$$

7. WEYL ASYMPTOTICS

In this section we obtain a Weyl-type asymptotic for the spectral shift function $\xi(\lambda, h)$ associated to operators H_0 and $H_1 = H_0 + V$. Here we assume that V is an electro-magnetic potential (8),

$$H_1 = - \sum_{j=1}^3 \alpha_j (ich\partial_j + eA_j) + mc^2\beta + ev.$$

In the following, we fix $I_0 \subset \mathbb{R} \setminus \{\pm mc^2\}$ and choose W_0 an open simply connected, relatively compact subset of Ω satisfying the assumption $(\mathbf{A}_\Omega^+)$ such that $I_0 = W_0 \cap \mathbb{R}$.

For the h -pseudo-differential and functional calculus for the Dirac operator, we refer to ([12],[10],[34],[19]). We recall that $H_\nu = Op_h^\omega(\mathcal{D}_\nu)$ and $\varphi(H_\nu)$ are h -pseudo-differential operators for a smooth function φ . The semi-classical symbol $\mathcal{D}_\nu$ is defined in (10).

Let us introduce the intervals $I_1, I_2 \subset I_0$ neighborhoods of λ_1, λ respectively such that, each $\lambda \in I_1 \cup I_2$ is a noncritical energy level for H (see Definition 2). Let $\varphi_j \in C_0^\infty(\mathbb{R}, \mathbb{R}^+)$ be such that

$$(46) \quad \varphi_1 = 1 \text{ on } I_1, \quad \varphi_2 = 1 \text{ on } I_2 \text{ and } \quad \varphi_1 + \varphi_2 + \varphi_3 = 1 \text{ on } I_0.$$

Consider a function $\theta(t) \in C_0^\infty(]-\delta_1, \delta_1[)$, $\theta(0) = 1$, $\theta(-t) = \theta(t)$, so that the Fourier transform $\hat{\theta}$ of θ satisfies $\hat{\theta}(\lambda) \geq 0$ on $\mathbb{R}$, and assume that there exist $0 < \epsilon_0 < 1$, $\delta_0 > 0$, such that $\hat{\theta}(\lambda) \geq \delta_0 > 0$ for $|\lambda| \leq \epsilon_0$. Next, we introduce

$$(\mathcal{F}_h^{-1}\theta)(\lambda) = (2\pi h)^{-1} \int e^{it\lambda h^{-1}} \theta(t) dt = (2\pi h)^{-1} \hat{\theta}(-h^{-1}\lambda).$$

To prove Theorem 3, we need the proposition:

Proposition 10. *For the trace involving H_ν , $\nu = 0, 1$, we have for $\lambda \in I_j$,*

$$(47) \quad \text{tr} \left(\left[(\mathcal{F}_h^{-1}\theta)(\lambda - H) \varphi_j(H) \right]_0^1 \right) = w_j(\lambda) h^{-3} + O(h^{-2}), \quad j = 1, 2,$$

with $w_j(\lambda) \in C_0^\infty(I_j)$ and $O(h^{-2})$ uniform with respect to $\lambda \in I_j$.

Proof. Proposition 10 is closed to the calculation of the trace in [5, Section 4] and to the appendix of [8] for the Schrödinger operator. But, here we use a trick of Robert [10]. We fix $j = 2$ (it is similar for $j = 1$). The proof of (47) is obtained following these two steps:

- First, we recall that $\lambda \in I_2$ and $\text{Supp } \theta(t) \subset [-\delta_1, \delta_1]$. Let us write

$$\begin{aligned} \mathcal{T} &= \text{tr} \left[(\mathcal{F}_h^{-1}\theta)(\lambda - H) \varphi_2(H) \right]_0^1 = \text{tr} \left[\int_0^1 \frac{\theta(t)}{2\pi h} e^{it(\lambda - H)h^{-1}} \varphi_2(H) dt \right]_0^1 \\ &= \frac{1}{2\pi h} \int e^{it\lambda h^{-1}} \theta(t) \text{tr} [e^{-itH \cdot h^{-1}} \varphi_2(H)]_0^1 dt \end{aligned}$$

In the order to calculate the trace

$$\text{tr}(f(H_1) - f(H_0)), \quad \text{for all } f \in C_0^\infty(\mathbb{R} \setminus \{\pm mc^2\})$$

we use [10, Proposition 3.2]. If we note $W(h) = Q - \frac{1}{2}i[Q, \mathcal{A}(h)]$ with $Q = H_1^2 - H_0^2$, $\mathcal{A}(h) = \frac{1}{2}(x \cdot h\partial_x + h\partial_x \cdot x)$ and $[Q, \mathcal{A}(h)] = Q\mathcal{A}(h) - \mathcal{A}(h)Q$, we have

$$(48) \quad \text{tr}(f(H_1) - f(H_0)) = \text{tr}(W(h)(H_1^2 - m^2c^4)^{-1}f(H_1)).$$

Applying the formula (48) for $f(\lambda) = e^{-it\lambda h^{-1}}\varphi_2(\lambda)$, we have

$$\mathcal{T} = \frac{1}{2\pi h} \int e^{it\lambda h^{-1}} \theta(t) \operatorname{tr} \left(W(h)(H_1^2 - m^2 c^4)^{-1} e^{-itH_1 h^{-1}} \varphi_2(H_1) \right) dt.$$

Remark 9. Of course $(H_1^2 - m^2 c^4)^{-1}$ is not well defined, however for $f \in C_0^\infty(\mathbb{R} \setminus \{\pm mc^2\})$, we can define $(H_1^2 - m^2 c^4)^{-1} f(H_1)$ as the self-adjoint operator $\varphi(H_1)$ where $\varphi \in C_0^\infty(\mathbb{R})$ satisfies:

$$\varphi(\lambda) = \begin{cases} (\lambda^2 - m^2 c^4)^{-1} f(\lambda) & \text{for } \lambda \neq \pm mc^2, \\ 0 & \text{for } \lambda = \pm mc^2. \end{cases}$$

• Now, we treat $\mathcal{T}$ following the analysis of [5, Section 4.2]. By h -pseudo-differential calculus, we obtain the existence of a h -pseudo-differential operator S which is trace class with symbol

$$(49) \quad s(x, y, \xi, h) \in \mathcal{S}^0(\langle x \rangle^{-\delta} \langle \xi \rangle^{-N}), \quad \forall N \in \mathbb{N}, \delta > 3,$$

having compact support in ξ and in $(x - y)$ (i.e. $\operatorname{supp}_{(x-y)}(s) = \{x - y, \exists \xi; (x, y, \xi, h) \in \operatorname{supp}(s)\}$ is compact) and support in $\{(x, \xi); |x| > R, (x, \xi) \in \mathcal{D}_1^{-1}(I_2)\}$, with $\mathcal{D}_1$ the semi-classical symbol of H_1 , so that

$$\mathcal{T} = \frac{1}{2\pi h} \operatorname{tr} \left(\int e^{it\lambda h^{-1}} \theta(t) e^{-itH_1 h^{-1}} S dt \right) + O(h^\infty).$$

Using Theorem 9 in Appendix A and the hypothesis of S by composition of Fourier integral operators, we obtain a Fourier integral operator $\tilde{\mathcal{U}}_t = \tilde{\mathcal{U}}_t^+ + \tilde{\mathcal{U}}_t^-$, such that for $|t| \leq \delta_1$ and δ_1 sufficiently small, we have

$$(50) \quad \|\tilde{\mathcal{U}}_t - e^{-itH_1 h^{-1}} S\|_{tr} = O(h^\infty),$$

where the kernel of the operator $\int e^{it\lambda h^{-1}} \theta(t) \tilde{\mathcal{U}}_t dt$ is equal to $\tilde{K}^+(x, y; h) + \tilde{K}^-(x, y; h)$ with

$$\tilde{K}^\pm(x, y; h) = \frac{1}{(2\pi h)^3} \iint e^{i(t\lambda + \Phi^\pm(t, x, \xi) - y \cdot \xi) h^{-1}} \theta(t) \tilde{E}^\pm(t, x, y, \xi; h) dt d\xi.$$

The amplitudes $\tilde{E}^\pm$, satisfy

$$\tilde{E}^\pm(t, x, y, \xi; h) \in \mathcal{S}^0(\langle x \rangle^{-\delta} \langle \xi \rangle^{-N}), \quad \forall N \in \mathbb{N}$$

and is compactly supported in ξ and in $(x - y)$.

Using the Taylor sum formula for the functions $\Phi^\pm(t, x, \xi)$ in a neighborhood of $t = 0$, we have:

$$\Phi^\pm(t, x, \xi) = x \cdot \xi - tH_1^\pm(x, \xi) + O(t^2).$$

We will deduce that $\mathcal{T} = \mathcal{T}^+ + \mathcal{T}^-$, with

$$\mathcal{T}^\pm = \frac{1}{(2\pi h)^4} \iiint e^{i(t\lambda + \Phi^\pm(t, x, \xi) - x \cdot \xi) h^{-1}} \theta(t) \tilde{E}^\pm(t, x, x, \xi; h) dt dx d\xi + O(h^\infty).$$

Moreover, the symbol $\tilde{E}^\pm(t, x, x, \xi; h)$ has support in $\{(x, \xi); |x| > R, |\xi| \leq C_1, (x, \xi) \in \mathcal{D}_1^{-1}(I_2)\}$, so that for all α and $|t| \leq \delta_1$, we have

$$(51) \quad |\partial^\alpha \tilde{E}^\pm(t, x, x, \xi; h)| \leq C_\alpha \langle x \rangle^{-\delta}, \quad \delta > 3.$$

The last estimate enables us to calculate $\mathcal{T}$ by using an infinite partition of unity

$$\sum_{\alpha \in \mathbb{N}^3} \Psi(x - \alpha) = 1, \quad \forall x \in \mathbb{R}^3,$$

where $\Psi \in C_0^\infty(K)$, $\Psi \geq 0$, K being a neighborhood of the unit cube. Consequently, for every fixed $h \in]0, h_0]$, we have

$$\begin{aligned} \mathcal{T}^\pm &= \frac{1}{(2\pi h)^4} \lim_{m \rightarrow \infty} \iiint e^{i(t\lambda + \Phi^\pm(t, x, \xi) - x \cdot \xi)h^{-1}} \theta(t) \\ &\quad \times \sum_{|\alpha| \leq m} \Psi(x - \alpha) \tilde{E}^\pm(t, x, x, \xi; h) dt dx d\xi + O(h^\infty) = \lim_{m \rightarrow \infty} I_m^\pm + O(h^\infty), \end{aligned}$$

and we reduce the problem to the analysis of the integrals $I_m^\pm$. Concerning the phase function, we observe that

$$(52) \quad t\lambda + \Phi^\pm(t, x, \xi) - x \cdot \xi = t(\lambda - H_1^\pm(x, \xi) + O(t)),$$

where $O(t)$ and its derivatives are uniformly bounded on the support of $\theta(t)\tilde{E}^\pm(t, x, x, \xi; h)$ since the derivatives of $(\Phi^\pm(t, x, \xi) - x \cdot \xi)$ are bounded on this set.

Now we look for critical points of the phase function $(t\lambda + \Phi^\pm(t, x, \xi) - x \cdot \xi)$. Putting the derivative with respect to t equal to 0, we see that $H_1^\pm(x, \xi) = \lambda + O(t)$. Since $\partial_{x, \xi} H_1^\pm(x, \xi) \neq 0$, when $H_1^\pm(x, \xi) = \lambda$, and putting the derivative of the phase function $t(\lambda - H_1^\pm(x, \xi) + O(t))$ with respect to $H_1^\pm(x, \xi)$ equal to 0, we have

$$t = O(t^2).$$

Then the phase is critical for $|t|$ small precisely when $t = 0$, $\lambda = H_1^\pm$. Near any such critical point we choose local coordinates $t, H_1^\pm(x, \xi), w_1, \dots, w_5$ and consider the Hessian of (52) with respect to $t, H_1^\pm(x, \xi)$ at the critical point:

$$\begin{pmatrix} \star & -1 \\ -1 & 0 \end{pmatrix}$$

This is a non-degenerate matrix of determinant -1 and of signature 0. By the stationary phase method we obtain

$$I_m^\pm = \frac{\psi^\pm(\lambda)}{(2\pi h)^3} \int_{\lambda = H_1^\pm} \sum_{|\alpha| \leq m} \Psi(x - \alpha) \tilde{E}^\pm(0, x, \xi, \lambda; h) L_\lambda^\pm(dw) + O(h^2),$$

where $L_\lambda^\pm(dw)$ is the Liouville measure on $\lambda = H_1^\pm$ and the remainder $O(h^{-2})$ is uniform with respect to $\lambda \in I_2$ and $m \in \mathbb{N}$. Here $\psi^\pm(\lambda) \in C_0^\infty(I_2)$. Taking the limit $\lim_{m \rightarrow \infty} I_m^\pm$, we obtain an asymptotics of $\mathcal{T}$. $\square$

Lemma 9. *With the above definitions of $\theta(t)$, $\xi(\lambda, h)$, $\varphi_j(\lambda)$, I_j , $j = 1, 2$, we have*

$$(53) \quad \int_{-\infty}^{\lambda} \mathcal{F}_h^{-1} \theta * \varphi_j \xi'(\mu, h) d\mu - \int_{-\infty}^{\lambda} \varphi_j(\mu) \xi'(\mu) d\mu = O(h^{-2}), \quad \lambda \in I_j.$$

Proof. We deal only with the analysis of (53) for $j = 2$ since that of $j = 1$ is similar. According to Theorem 2, there exists a holomorphic function $r(z, h)$ in Ω such that for all $\lambda \in I_0 = W_0 \cap \mathbb{R}$, we have

$$\xi'(\lambda, h) = \frac{1}{\pi} \operatorname{Im} r(\lambda, h) + \sum_{\substack{w \in \operatorname{Res}(H_1) \cap \Omega \\ \operatorname{Im} w \neq 0}} \frac{-\operatorname{Im} w}{\pi |\lambda - w|^2} + \sum_{w \in \operatorname{Res}(H_1) \cap I_0} \delta_w(\lambda),$$

where $r(z, h)$ satisfies the following estimate:

$$(54) \quad |r(z, h)| \leq C(W)h^{-3}, \quad z \in W,$$

with $C(W) > 0$ independent of h . Let us denote

$$(55) \quad M_{\varphi_2}(\lambda) = \sum_{\substack{w \in \text{Res}(H_1) \cap \Omega \\ \text{Im} w \neq 0}} \int_{-\infty}^{\lambda} \frac{-\text{Im } w}{\pi |\lambda - w|^2} \varphi_2(\mu) d\mu + \sum_{w \in \text{Res}(H_1) \cap]c_0, \lambda]} \varphi_2(w).$$

Using the Cauchy inequalities and (54), it follows easily that

$$G'_{\varphi_2}(\lambda) = O(h^{-3}) \text{ and } G''_{\varphi_2}(\lambda) = O(h^{-3}),$$

and we immediately obtain

$$(56) \quad \mathcal{F}_h^{-1} \theta * G'_{\varphi_2} - G'_{\varphi_2} = O(h^{-2}).$$

Now, we want to apply a Tauberian theorem (see [34, Theorem V-13]) for the increasing function $M_{\varphi_2}(\lambda)$. For this purpose, we need the estimates

$$(57) \quad M_{\varphi_2}(\lambda) = O(h^{-3}), \quad \frac{d}{d\lambda}(\mathcal{F}_h^{-1} \theta * M_{\varphi_2})(\lambda) = O(h^{-3}), \quad \forall \lambda \in \mathbb{R},$$

and the equality $M_{\varphi_2}(\mu) = G_{\varphi_2}(\mu) = 0, \quad \mu \leq \inf I_2$.

The first estimate in (57) follows easily from the equation (55) with the upper bound of the number of the resonances in Ω (see Theorem 1), and the second follows from (47) and the equation

$$\frac{d}{d\lambda}(\mathcal{F}_h^{-1} \theta * M_{\varphi_2})(\lambda) = \mathcal{F}_h^{-1} \theta * \varphi_2 \xi'(\lambda) - \frac{d}{d\lambda}(\mathcal{F}_h^{-1} \theta * G_{\varphi_2})(\lambda).$$

Then, according to the Tauberian theorem we have

$$(\mathcal{F}_h^{-1} \theta * M_{\varphi_2})(\lambda) = M_{\varphi_2}(\lambda) + O(h^{-2}),$$

this enabled us to obtain

$$\begin{aligned} \int_{-\infty}^{\lambda} \varphi_2(\mu) \xi'(\mu) d\mu &= M_{\varphi_2}(\lambda) + \int_{-\infty}^{\lambda} G'_{\varphi_2}(\mu) d\mu \\ &= \int_{-\infty}^{\lambda} \frac{d}{d\mu}(\mathcal{F}_h^{-1} \theta * M_{\varphi_2} + \mathcal{F}_h^{-1} \theta * G_{\varphi_2})(\mu) d\mu + O(h^{-2}) \\ &= \int_{-\infty}^{\lambda} \mathcal{F}_h^{-1} \theta * \varphi_2 \xi'(\mu, h) d\mu + O(h^{-2}). \end{aligned}$$

□

Proof of Theorem 3. For $\lambda_1 \in I_1$, $\lambda \in I_2$, using the functions defined in (46), we have

$$\begin{aligned}
 \xi(\lambda, h) - \xi(\lambda_1, h) &= \int_{-\infty}^{\lambda} \varphi_1(\mu) \xi'(\mu, h) d\mu - \int_{-\infty}^{\lambda_1} \varphi_2(\mu) \xi'(\mu, h) d\mu \\
 &\quad - \int_{-\infty}^{\lambda_1} \varphi_1(\mu) \xi'(\mu, h) d\mu + \int_{-\infty}^{\lambda} \varphi_2(\mu) \xi'(\mu, h) d\mu \\
 &\quad + \int_{\lambda_1}^{\lambda} \varphi_3(\mu) \xi'(\mu, h) d\mu.
 \end{aligned}
 \tag{58}$$

Since $\varphi_j = 0$ on I_{3-j} for $j = 1, 2$, the first term (resp. the second term) is independent of $\lambda \in I_2$ (resp. $\lambda_1 \in I_1$) and is equal to $\text{tr}[\varphi_1(H.)]_0^1 = C(\varphi_1)h^{-3} + O(h^{-2})$ (resp. $\text{tr}[\varphi_2(H.)]_0^1 = C(\varphi_2)h^{-3} + O(h^{-2})$), where $C(\varphi_j)$ is a constant depending on φ_j for $j = 1, 2$. Since $\varphi_3 = 0$ on I_j , $j = 1, 2$, the last term is independent of $\lambda \in I_2$, $\lambda_1 \in I_1$ and is equal to $C(\varphi_3)h^{-3} + O(h^{-2})$, where $C(\varphi_3)$ is a constant depending on φ_3 . The proof of this results is based on the functional calculus in the framework of h -pseudo-differential operators.

Using the equations (47), (53) and (58) we complete the proof of the asymptotic expansion (13) by writing

$$\begin{aligned}
 (\mathcal{F}_h^{-1}\theta * (\varphi_j \xi'))(\lambda) &= \langle (\mathcal{F}_h^{-1}\theta)(\lambda - \cdot) \varphi_j(\cdot), \xi' \rangle \\
 &= \text{tr} \left(\left[(\mathcal{F}_h^{-1}\theta)(\lambda - H.) \varphi_j(H.) \right]_0^1 \right) \\
 &= w_j(\lambda)h^{-3} + O(h^{-2}), \quad j = 1, 2.
 \end{aligned}
 \tag{59}$$

It remains to compute the Weyl term (14).

According to the definition of the spectral shift function $\xi(\lambda, h)$ in (5), we have:

$$\langle \xi'(\lambda, h), \varphi(\lambda) \rangle = \text{tr}(\varphi(H_1) - \varphi(H_0)), \quad \varphi(\lambda) \in C_0^\infty(\mathbb{R}).
 \tag{60}$$

We use weak asymptotics which is a direct consequence of functional calculus in the framework of h -pseudo-differential operators, as settled in [12], [34], [10]. We find

$$H_\nu = Op_h^\omega(\mathcal{D}_\nu), \quad \nu = 0, 1, \quad (\mathcal{D}_\nu \text{ defined in (10)}),$$

and

$$\begin{aligned}
 \text{tr}(\varphi(H_1) - \varphi(H_0)) &= h^{-3} \sum_{j \geq 0} \gamma_j(\varphi) h^j \\
 &= h^{-3} \gamma_0(\varphi) + O(h^{-2}),
 \end{aligned}$$

with $\gamma_0(\varphi) = (2\pi)^{-3} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \text{tr}(\varphi(\mathcal{D}_1(x, \xi)) - \varphi(\mathcal{D}_0(x, \xi))) dx d\xi$.
($\text{tr}(A)$ is the trace of the matrix A).

The matrix $\mathcal{D}_\nu(x, \xi)$ is Hermitian and has two eigenvalues $H_\nu^\pm(x, \xi)$ (see (11)), then

$$\text{tr}(\varphi(\mathcal{D}_1) - \varphi(\mathcal{D}_0)) = 2(\varphi(H_1^+) + \varphi(H_1^-) - \varphi(H_0^+) - \varphi(H_0^-)).$$

According to the asymptotic expansions (13) and (60) we obtain

$$w(\lambda, \lambda_1) = w(\lambda) - w(\lambda_1),$$

with

$$w(\lambda) = \frac{1}{4\pi^3} \int_{\mathbb{R}^3} \left(\int_{H_1^+(x,\xi) \leq \lambda} d\xi - \int_{H_0^+(x,\xi) \leq \lambda} d\xi - \int_{H_1^-(x,\xi) \geq \lambda} d\xi + \int_{H_0^-(x,\xi) \geq \lambda} d\xi \right) dx.$$

Putting $\zeta_\nu = c\xi - \nu eA(x)$ for $\nu = 0, 1$ and $\zeta = r\omega$ ($\omega \in S^2$), we get

$$\begin{aligned} \pm H_\nu^\pm \leq \pm \lambda &\Leftrightarrow \left(\zeta_\nu^2 + (mc^2 + \nu \frac{e(v_+ - v_-)}{2})^2 \right)^{\frac{1}{2}} \pm \nu \frac{e(v_+ + v_-)}{2} \leq \pm \lambda \\ &\Leftrightarrow \left(\zeta_\nu^2 + (mc^2 + \nu \frac{e(v_+ - v_-)}{2})^2 \right)^{\frac{1}{2}} \leq \pm \left(\lambda - \nu \frac{e(v_+ + v_-)}{2} \right), \end{aligned}$$

thus

$$\begin{aligned} \int_{H_1^+(x,\xi) \leq \lambda} d\xi - \int_{H_1^-(x,\xi) \geq \lambda} d\xi &= \frac{4\pi}{3} \left(\left(\lambda - \frac{e(v_+ + v_-)}{2} \right)_+^2 - (mc^2 + \frac{e(v_+ - v_-)}{2})^2 \right)_+^{\frac{3}{2}} \\ &\quad - \frac{4\pi}{3} \left(\left(\lambda - \frac{e(v_+ + v_-)}{2} \right)_-^2 - (mc^2 + \frac{e(v_+ - v_-)}{2})^2 \right)_+^{\frac{3}{2}}, \end{aligned}$$

and

$$- \int_{H_0^+(x,\xi) \leq \lambda} d\xi + \int_{H_0^-(x,\xi) \geq \lambda} d\xi = \mp \frac{4\pi}{3} (\lambda^2 - (mc^2)^2)_+^{\frac{3}{2}}, \quad \text{for } \pm \lambda \geq 0,$$

with $(x)_+ = \max(x, 0)$ and $(x)_- = \max(-x, 0)$ for $x \in \mathbb{R}$. $\square$

Remark 10. Theorem 3 can be extended to the pair operators $(H_1 = H_0 + V_1, H_2 = H_0 + V_2)$, where the potentials V_1, V_2 are electro-magnetic potentials

$$V(x) = e(-\alpha \cdot A + v)(x) = - \sum_{j=1}^3 \alpha_j \cdot eA_j(x) + e \begin{pmatrix} v_+(x)I_2 & 0 \\ 0 & v_-(x)I_2 \end{pmatrix}$$

satisfying the assumption $(\mathbf{A}_V)$ with $\delta > 0$ (or $\|V_j(x)\| \rightarrow 0$) and the potential $V = V_2 - V_1$ satisfies the assumption $(\mathbf{A}_V)$ with $\delta > 3$:

For all λ, λ_1 noncritical energy levels for H_1, H_2 such that $\pm mc^2 \notin]\lambda_1, \lambda[$ and $h \in]0, h_0[$, we have the asymptotic expansion

$$(61) \quad \xi(\lambda, h) - \xi(\lambda_1, h) = w(\lambda, \lambda_1)h^{-3} + O(h^{-2}).$$

Here the $O(h^{-2})$ is uniform for λ (resp. λ_1) in a small interval I_2 (resp. I_1). The first term $w(\lambda, \lambda_1) \in C^\infty(I_2 \times I_1)$ is given by

$$w(\lambda, \lambda_1) = w(\lambda) - w(\lambda_1)$$

with,

$$(62) \quad w(\lambda) = \frac{1}{3\pi^2} \int_{\mathbb{R}^3} [W_+(\lambda, v_+, v_-) - W_+(\lambda, v_+, v_-)]_1^2 dx$$

$$\text{where } W_\pm(\lambda, a, b) = \left(\left(\lambda - \frac{e(a+b)}{2} \right)_\pm^2 - \left(mc^2 + \frac{e(a-b)}{2} \right)^2 \right)_+^{\frac{3}{2}}.$$

In this setting, we does not have a formula like (48). But it could be possible to use the approach to Bruneau-Petkov in [8]. For that we need more informations on the approximation of the propagator $e^{-itH_j h^{-1}}$ by the Fourier integral operator $\mathcal{U}_t$.

8. RESONANCES IN SMALL DOMAINS

In this section, we assume that the hamiltonian $H = H_0 + V$, where V is an electro-magnetic potential:

$$H = - \sum_{j=1}^3 \alpha_j (i\hbar \partial_j + eA_j) + mc^2 \beta + ev.$$

8.1. Upper bound for the number of resonances in domains of width h . We adapt, for the Dirac operator, Theorem 1 of [9] which rests on a representation formula for the spectral shift function (see Theorem 2).

Theorem 7. *Suppose that each $\lambda \in [E_0, E_1]$ is a non-critical energy level for H . Then for $h \in]0, h_0]$, the following assertions are equivalents:*

- (i) *There exist positive constants B, C, b, h_0 , such that for any $\lambda \in [E_0 - b, E + b]$, $h \in]0, h_0]$ and $h/B \leq \rho \leq B$, we have*

$$\#\{z \in \mathbb{C} : z \in \text{Res}(H), |z - \lambda| \leq \rho\} \leq C\rho h^{-3}.$$

- (ii) *There exist positive constants $B_1, C_1, \varepsilon_1, h_1$, such that for any $\lambda \in [E_0 - \varepsilon_1, E_1 + \varepsilon_1]$, $h \in]0, h_1]$ and $h/B_1 \leq \rho \leq B_1$, we have*

$$|\xi(\lambda + \rho, h) - \xi(\lambda - \rho, h)| \leq C\rho h^{-3}.$$

As a consequence of Theorem 7, we have an upper bound $O(h^{-2})$ for the number of the resonances for the semi-classical Dirac operator close to a non-critical energy level in a domain of width h :

Proposition 11. *Assume that V is the electro-magnetic potential (8) satisfying the assumption $(\mathbf{A}_V)$ with $\delta > 3$. We suppose also that each $\lambda \in [E_0, E_1]$ is a non-critical energy level for H . There are positive constants C, B, b, h_0 such that for any $\lambda \in [E_0 - b, E + b]$, $h \in]0, h_0]$ and $h/B \leq \rho \leq B$, we have*

$$\#\{z \in \mathbb{C} : z \in \text{Res}(H), |z - \lambda| \leq \rho\} \leq C\rho h^{-3}.$$

Proof. It follows from Theorem 7 and equation (13). $\square$

8.2. Breit-Wigner approximation. In this part, we consider small domains of width h , and we prove a Breit-Wigner approximation for $\xi(\lambda, h)$ (see [31], [32], [16], [6], [8]). Let $\eta(\lambda, h)$ be the real analytic function defined by

$$\eta(\lambda, h) = \xi(\lambda, h) - \#\{\mu \in [E_0, \lambda] : \mu \in \sigma_d(H)\}.$$

Using Proposition 11 and the arguments used in [8, Section 6], we obtain a Breit-Wigner approximation for the derivative of the spectral shift function $\xi(\lambda, h)$.

Theorem 8. (Breit-Wigner) *Assume that V is an electro-magnetic potential (8), for any $\lambda \in [E_0, E_1]$ non-critical energy level for H , $0 < \rho < h/B$, $0 < B_1 < B$, and h sufficiently small, we have*

$$\eta(\lambda + \rho, h) - \eta(\lambda - \rho, h) = \sum_{\substack{w \in \text{Res}(H) \\ \text{Im } w \neq 0, |w - \lambda| < h/B_1}} \omega_{\mathbb{C}_-}(w, [\lambda - \rho, \lambda + \rho]) + O(\rho)h^{-3},$$

where $B > 0$ is a positive constant and $\omega_{\mathbb{C}_-}$ is the harmonic measure

$$\omega_{\mathbb{C}_-}(w, E) = -\frac{1}{\pi} \int_E \frac{\text{Im}(w)}{|t - w|^2} dt, \quad E \subset \mathbb{R} = \partial\mathbb{C}_-.$$

Using Theorem 7 and repeating with little modifications the arguments used in [7, Section 6], we obtain the following corollary which implies also trace formula in small domains.

Corollary 1. *Under the assumptions of Theorem 8 and supposing that $[E_0, E_1]$ is a non-critical energy level for H , for each $E \in [E_0, E_1]$ there exist constants $C_2 > C_1 > 0$, $h_0 > 0$ so that for $|\lambda - E| \leq C_1 h$, $h \in]0, h_0]$, we have*

$$(63) \quad \xi'(\lambda, h) = -\frac{1}{\pi} \sum_{\substack{w \in \text{Res}(H) \\ |E - w| \leq C_2 h}} \frac{\text{Im}(w)}{|\lambda - w|^2} + \sum_{\substack{w \in \sigma_d(H) \\ |E - w| \leq C_1 h}} \delta_w(\lambda) + O(h^{-3}).$$

Here $\delta_w(\cdot)$ is the Dirac mass at $w \in \mathbb{R}$.

Appendix A. Construction of $\mathcal{U}_t$

In this appendix, we construct a parametrix at small time of the propagator of the Dirac equation in an external electro-magnetic field

$$ih\partial_t\psi = H_1\psi,$$

with $H_1 = H_0 + V$. Here H_0 is the selfadjoint operator defined in (1) and V is an electro-magnetic potential (8).

Theorem 9. (Approximation of the propagator) *There exist $\delta_1 > 0$ small enough and a Fourier integral operator $\mathcal{U}_t = \mathcal{U}_t^+ + \mathcal{U}_t^-$ with*

$$\mathcal{U}_t^\pm f(y) = \frac{1}{(2\pi h)^3} \iint e^{i(\Phi^\pm(t, x, \xi) - y \cdot \xi)h^{-1}} E^\pm(t, x, y, \xi; h) f(y) d\xi dy,$$

defined for $|t| < \delta_1$ such that:

- The amplitude $E^\pm(t, x, y, \xi; h) \in \mathcal{S}^0(1)$.
- $\|\mathcal{U}_t - e^{-itH_1h^{-1}}\| = O(h^\infty)$, uniformly for $|t| < \delta_1$.
- The phase function $\Phi^\pm(t, x, \xi) - x \cdot \xi$ and their derivatives $\partial_t^\alpha \partial_x^\beta \partial_\xi^\gamma (\Phi^\pm(t, x, \xi) - x \cdot \xi)$ are uniformly bounded for $(t, x, \xi) \in [-\delta_1, \delta_1] \times \mathbb{R}^3 \times B(0, C_1)$, $(\alpha, \beta, \gamma) \neq (0, 0, 0)$ and $C_1 > 0$ (see (68)).

With a different approach, such result has been obtained by Yajima [46] for a scalar electric potential ($v_+ = v_-$).

Proof. We consider the equivalent problem for $\mathcal{U}_t$

$$(64) \quad \begin{cases} ih\partial_t \mathcal{U}_t - H_1 \mathcal{U}_t = 0, \\ \mathcal{U}_0 = I. \end{cases}$$

We solve this problem using the B.K.W. method. We assume that the kernel of the operator $\mathcal{U}_t$ is K_t , where

$$K_t(x, y; h) = \frac{1}{(2\pi h)^3} \int e^{i(\Phi(t, x, \xi) - y \cdot \xi)h^{-1}} E(t, x, y, \xi; h) d\xi,$$

with $E(t, x, y, \xi; h) = E_0(t, x, y, \xi) + hE_1(t, x, y, \xi) + \dots$.

Thus, if we look for $E(t, x, y, \xi; h)$ having the asymptotic expansion above, it is enough to solve (in some fixed neighborhood of $t = 0$) the sequence of equations

$$(65) \quad \begin{cases} 0 &= (\partial_t \Phi(t, x, \xi) + c\alpha \cdot \nabla_x \Phi - e\alpha \cdot A + mc^2\beta + ev) E_0, \\ i(\partial_t + c\alpha \cdot \nabla_x) E_j &= (\partial_t \Phi(t, x, \xi) + c\alpha \cdot \nabla_x \Phi - e\alpha \cdot A + mc^2\beta + ev) E_{j+1}, \\ E_0(0, x, \xi) &= I_4, \\ E_j(0, x, \xi) &= 0, \text{ for } j \geq 1. \end{cases}$$

On the support of E_0 , we deduce the eikonal equation

$$(66) \quad \begin{cases} \det(\partial_t \Phi(t, x, \xi) + c\alpha \cdot \nabla_x \Phi - e\alpha \cdot A + mc^2\beta + ev) = 0, \\ \Phi(0, x, \xi) = x \cdot \xi. \end{cases}$$

The system (66) is equivalent to

$$(67) \quad \begin{cases} \partial_t \Phi^\pm(t, x, \xi) + H_1^\pm(x, \nabla_x \Phi) = 0, \quad (\text{see (11)}), \\ \Phi^\pm(0, x, \xi) = x \cdot \xi. \end{cases}$$

The last system can be solved by Hamilton-Jacobi method (see [2]) and all derivatives

$$(68) \quad \partial_t^\alpha \partial_x^\beta \partial_\xi^\gamma (\Phi^\pm(t, x, \xi) - x \cdot \xi)$$

are uniformly bounded for $(t, x, \xi) \in [-\delta_1, \delta_1] \times \mathbb{R}^3 \times B(0, C_1)$ and $(\alpha, \beta, \gamma) \neq (0, 0, 0)$. Using the Taylor formula in a neighborhood of $t = 0$, the two solutions of (67) satisfy:

$$\Phi^\pm(t, x, \xi) = x \cdot \xi - tH_1^\pm(x, \xi) + O(t^2).$$

Then $\mathcal{U}_t = \mathcal{U}_t^+ + \mathcal{U}_t^-$, and the kernel of the operator $\mathcal{U}_t$ is $K_t = K_t^+ + K_t^-$, with

$$K_t^\pm(x, y; h) = \frac{1}{(2\pi h)^3} \int e^{i(\Phi^\pm(t, x, \xi) - y \cdot \xi)h^{-1}} E^\pm(t, x, y, \xi; h) d\xi.$$

We look for the amplitude $E^\pm(t, x, y, \xi; h)$ having an asymptotic expansions according to the power of h :

$$E_0^\pm(t, x, y, \xi) + hE_1^\pm(t, x, y, \xi) + \dots$$

Consequently, the coefficients $E_j^\pm(t, x, y, \xi)$ are the solutions of the transport equations

$$(69) \quad \begin{cases} 0 &= (\partial_t \Phi^\pm + c\alpha \cdot \nabla_x \Phi^\pm - e\alpha \cdot A + mc^2\beta + ev) E_0^\pm, \\ i(\partial_t + c\alpha \cdot \nabla_x) E_j^\pm &= (\partial_t \Phi^\pm + c\alpha \cdot \nabla_x \Phi^\pm - e\alpha \cdot A + mc^2\beta + ev) E_{j+1}^\pm, \\ E_j^+(0, x, \xi) + E_j^-(0, x, \xi) &= 0 \text{ for } j \geq 1, \\ E_0^\pm(0, x, \xi) &= \Pi_1^\pm(x, \xi), \end{cases}$$

with $\Pi_1^\pm(x, \xi)$ defined by (12).

Resolution of (69).

Let us denote by $L = \partial_t + c\alpha \cdot \nabla_x$, with $\alpha \cdot \nabla_x = \sum_{j=1}^3 \alpha_j \partial_{x_j}$. The matrix

$$\mathcal{M}^\pm = \partial_t \Phi^\pm + c\alpha \cdot \nabla_x \Phi^\pm - e\alpha \cdot A + mc^2\beta + ev,$$

is Hermitian and has two real eigenvalues linearly independent with multiplicity 2.

First, we multiply the system (69) by the column-vector $N_1 = (1, 0, 0, 0)^\dagger$, the subscript $\dagger$ designates the complex conjugate of the transpose. We denote

$$(70) \quad E_{j,1}^\pm = E_j^\pm N_1 \text{ for } j = 1, 2, \dots, \quad E_{0,1}^\pm(0, x, \xi) = \Pi_1^\pm(x, \xi) N_1.$$

Since $\det(\mathcal{M}^\pm) = 0$, there exist $l_k^\pm$ and $r_k^\pm$, left and right eigenvectors of the matrix $\mathcal{M}^\pm$, corresponding to the eigenvalue zero, such that

$$(71) \quad \mathcal{M}^\pm r_k^\pm = 0, \quad l_k^\pm \mathcal{M}^\pm = 0, \quad l_k^\pm = (r_k^\pm)^\dagger, \quad k = 1, 2,$$

(here $r_k^\pm$ is a column-vector and $l_k^\pm$ is a row-vector). We choose

$$r_1^\pm = \begin{pmatrix} u^+ \\ 0 \\ v^+ \\ w_+^\pm \end{pmatrix}, \quad r_2^\pm = \begin{pmatrix} 0 \\ u^+ \\ w_-^\pm \\ -v^+ \end{pmatrix}, \quad r_1^\mp = \begin{pmatrix} w_+^\mp \\ v^- \\ 0 \\ u^- \end{pmatrix}, \quad r_2^\mp = \begin{pmatrix} -v^- \\ w_-^\mp \\ u^- \\ 0 \end{pmatrix},$$

$$(72) \quad l_\nu^\pm r_k^\pm = (\mp 2p_5^\pm u^\pm) \delta_{\nu k}, \quad \nu, k = 1, 2.$$

Here $u^\pm, v^\pm$ and $w_\pm^\pm$ are defined by

$$u^\pm = p_4^\pm \mp p_5^\pm, \quad v^\pm = p_3^\pm, \quad w_+^\pm = \pm p_1^\pm + ip_2^\pm, \quad w_-^\pm = \pm p_1^\pm - ip_2^\pm,$$

where $p_4^\pm = mc^2 + \frac{e(v_+ - v_-)}{2}$, $p_5^\pm = \partial_t \Phi^\pm + \frac{e(v_+ + v_-)}{2}$, $p_j^\pm = c\partial_{x_j} \Phi^\pm - eA_j$, for $j = 1, 2, 3$.

It is easy to see that the vector-valued functions $r_k^\pm(t, x, \xi)$ and $l_k^\pm(t, x, \xi)$ can be chosen to be smooth in t and x and different from 0 everywhere. All the derivatives of $r_k^\pm, l_k^\pm$, $k = 1, 2$, are uniformly bounded for $(t, x, \xi) \in [-\delta_1, \delta_1] \times \mathbb{R}^3 \times B(0, C_1)$. Then it follows from the first equation in (69) that

$$E_{0,1}^\pm = \sigma_{0,1}^\pm(t, x, \xi) r_1^\pm(t, x, \xi) + \sigma_{0,2}^\pm(t, x, \xi) r_2^\pm(t, x, \xi),$$

where $\sigma_{0,1}^\pm, \sigma_{0,2}^\pm$ are scalar-valued functions. If we multiply the second equation in (69) for $j = 0$ on the left by $l_k^\pm$ for $k = 1, 2$, we deduce the following differential equations for $\sigma_{0,k}^\pm$:

$$\begin{cases} l_1^\pm L(\sigma_{0,1}^\pm r_1^\pm) + l_1^\pm L(\sigma_{0,2}^\pm r_2^\pm) = 0, \\ l_2^\pm L(\sigma_{0,1}^\pm r_1^\pm) + l_2^\pm L(\sigma_{0,2}^\pm r_2^\pm) = 0. \end{cases}$$

We conclude

$$(73) \quad \begin{cases} l_1^\pm r_1^\pm \partial_t(\sigma_{0,1}^\pm) + c \sum_{j=1}^3 l_1^\pm \alpha_j r_1^\pm \partial_{x_j}(\sigma_{0,1}^\pm) + c \sum_{j=1}^3 l_1^\pm \alpha_j r_2^\pm \partial_{x_j}(\sigma_{0,2}^\pm) \\ \quad + l_1^\pm L(r_1^\pm) \sigma_{0,1}^\pm + l_1^\pm L(r_2^\pm) \sigma_{0,2}^\pm = 0, \\ l_2^\pm r_2^\pm \partial_t(\sigma_{0,2}^\pm) + c \sum_{j=1}^3 l_2^\pm \alpha_j r_2^\pm \partial_{x_j}(\sigma_{0,2}^\pm) + c \sum_{j=1}^3 l_2^\pm \alpha_j r_1^\pm \partial_{x_j}(\sigma_{0,1}^\pm) \\ \quad + l_2^\pm L(r_1^\pm) \sigma_{0,1}^\pm + l_2^\pm L(r_2^\pm) \sigma_{0,2}^\pm = 0. \end{cases}$$

We now use Lemma 10 (see below) in the system (73). Since $p_5^\pm \neq 0$, $u^\pm = p_4^\pm \mp p_5^\pm \neq 0$ then, after multiplying (73) by $(\mp 2p_5^\pm u^\pm)^{-1}$, the system (73) can be written as

$$(74) \quad D^\pm \sigma_0^\pm = M^\pm \sigma_0^\pm := (\mp 2p_5^\pm u^\pm)^{-1} \begin{pmatrix} l_1^\pm L(r_1^\pm) & l_1^\pm L(r_2^\pm) \\ l_2^\pm L(r_1^\pm) & l_2^\pm L(r_2^\pm) \end{pmatrix} \begin{pmatrix} \sigma_{0,1}^\pm \\ \sigma_{0,2}^\pm \end{pmatrix},$$

with $D^\pm = \partial_t + a^\pm \cdot \nabla_x = \partial_t + \sum_{j=1}^3 a_j^\pm(t, x) \partial_{x_j}$, and

$$a^\pm = c(\mp 2p_5^\pm u^\pm)^{-1} (l_1^\pm \alpha_1 r_1^\pm, l_1^\pm \alpha_2 r_1^\pm, l_1^\pm \alpha_3 r_1^\pm).$$

Thus the function $\sigma_{0,k}^\pm$ can be found if its value is known for $t = 0$, and it is as smooth as $\sigma_{0,k}^\pm(0, x, \xi)$ (for more details, see a method for solving a similar equation in [37]). The equality

$$E_{0,1}^\pm(0, x, \xi) = \sigma_{0,1}^\pm(0, x, \xi) r_1^\pm(0, x, \xi) + \sigma_{0,2}^\pm(0, x, \xi) r_2^\pm(0, x, \xi) = \Pi_1^\pm(x, \xi) N_1,$$

gives the value of $\sigma_0^\pm$ at $t = 0$.

Since, all the derivatives of $\sigma_{0,k}^\pm$, $r_k^\pm$, for $k = 1, 2$, are uniformly bounded, then all the derivatives $(\partial_t^\alpha \partial_x^\beta \partial_\xi^\gamma E_{0,1}^\pm)$ are uniformly bounded for $(\alpha, \beta, \gamma) \in \mathbb{N} \times \mathbb{N}^3 \times \mathbb{N}^3$.

It follows from the second equation in (69) for $j = 0$, that

$$iLE_{0,1}^\pm = \mathcal{M}^\pm E_{1,1}^\pm,$$

i.e., $E_{1,1}^\pm = \sigma_{1,1}^\pm r_1^\pm + \sigma_{1,2}^\pm r_2^\pm + h_1^\pm$, where $\sigma_{1,k}^\pm$ is a scalar-valued function for $k = 1, 2$, and $h_1^\pm$ is expressed in term of $LE_{0,1}^\pm$. To find $\sigma_{1,k}^\pm$ it is sufficient to multiply the second equation in (69) for $j = 1$ on the left by $l_k^\pm$ for $k = 1, 2$. Then

$$\begin{cases} l_1^\pm L(\sigma_{1,1}^\pm r_1^\pm) + l_1^\pm L(\sigma_{1,2}^\pm r_2^\pm) + l_1^\pm L(h_1^\pm) = 0, \\ l_2^\pm L(\sigma_{1,1}^\pm r_1^\pm) + l_2^\pm L(\sigma_{1,2}^\pm r_2^\pm) + l_2^\pm L(h_1^\pm) = 0. \end{cases}$$

From this equation, $\sigma_{1,k}^\pm$ can be found if the function $\sigma_{1,k}(0, x)$ is known. With the same process, for all $j = 1, 2, \dots$, we obtain

$$\begin{cases} \sigma_{j,1}^\pm r_1^\pm + \sigma_{j,2}^\pm r_2^\pm + h_j^\pm = E_{j,1}^\pm, \\ l_1^\pm L(\sigma_{j,1}^\pm r_1^\pm) + l_1^\pm L(\sigma_{j,2}^\pm r_2^\pm) + l_1^\pm L(h_j^\pm) = 0, \\ l_2^\pm L(\sigma_{j,1}^\pm r_1^\pm) + l_2^\pm L(\sigma_{j,2}^\pm r_2^\pm) + l_2^\pm L(h_j^\pm) = 0. \end{cases}$$

For $t = 0$, $j = 1, 2, \dots$, we have

$$\sigma_{0,1}^\pm r_1^\pm + \sigma_{0,2}^\pm r_2^\pm = \Pi_1^\pm N_1, \quad \sigma_{j,1}^+ r_1^+ + \sigma_{j,1}^- r_1^- + \sigma_{j,2}^+ r_2^+ + \sigma_{j,2}^- r_2^- = -(h_j^+ + h_j^-),$$

and the quantity $h_j^\pm$ is determined if $E_{0,1}^\pm, E_{1,1}^\pm, \dots, E_{j-1,1}^\pm$, are known. Solving the differential equation for $\sigma_j^\pm = \begin{pmatrix} \sigma_{j,1}^\pm \\ \sigma_{j,2}^\pm \end{pmatrix}$, we find these functions for all sufficiently small t .

Repeating this block of calculus, multiplying by $N_2 = (0, 1, 0, 0)^\dagger$, $N_3 = (0, 0, 1, 0)^\dagger$ and $N_4 = (0, 0, 0, 1)^\dagger$ instead of N_1 in (70), we find $E_{j,2}^\pm = E_j^\pm N_2$, $E_{j,3}^\pm = E_j^\pm N_3$ and $E_{j,4}^\pm = E_j^\pm N_4$. Consequently, we have:

Proposition 12. *There exists a family of matrices*

$$E_j^\pm = (E_{j,1}^\pm, E_{j,2}^\pm, E_{j,3}^\pm, E_{j,4}^\pm), \quad \text{for } j \geq 0,$$

solution of (69). Moreover, for all $j \geq 0$, $E_j^\pm \in C^\infty$ and all derivatives $(\partial_t^\alpha \partial_x^\beta \partial_\xi^\gamma E_j^\pm)$ are uniformly bounded for all $(t, x, \xi) \in [-\delta_1, \delta_1] \times \mathbb{R}^3 \times B(0, C_1)$ and $(\alpha, \beta, \gamma) \in \mathbb{N} \times \mathbb{N}^3 \times \mathbb{N}^3$.

Consequently, Borel process provides a symbol $E^\pm(t, x, y, \xi; h) \in \mathcal{S}^0(1)$ with compact support on ξ and $(x - y)$ with $E_0^\pm(t, x, y, \xi) + hE_1^\pm(t, x, y, \xi) + \dots$ its asymptotic expansion.

Desired estimate.

Next, we remark that for all $N \in \mathbb{N}$:

$$(75) \quad \begin{aligned} (ih\partial_t - H_1) \left(e^{i(\Phi^\pm(t, x, \xi) - y \cdot \xi)h^{-1}} \sum_{j=0}^N h^j E_j^\pm \right) &= e^{i(\Phi^\pm(t, x, \xi) - y \cdot \xi)h^{-1}} \\ &\quad \times \sum_{j=0}^N (ihL(E_j^\pm) + \mathcal{M}^\pm E_j^\pm) h^j \\ &= P_N(t, x, \xi; h) h^N, \end{aligned}$$

and all derivatives $D_{x, \xi}^\alpha P_N(t, x, \xi; h)$ are bounded as $h \rightarrow 0$ for all α . Then for all $N \in \mathbb{N}$,

$$(76) \quad \begin{cases} ih\partial_t \mathcal{U}_t - H_1 \mathcal{U}_t &= O(h^N), \\ \mathcal{U}_0 &= I + O(h^N), \end{cases}$$

thus

$$(77) \quad \begin{cases} \frac{d}{dt}(e^{+itH_1} \mathcal{U}_t) &= O(h^N), \\ \mathcal{U}_0 &= I + O(h^N), \end{cases}$$

where $O(h^N)$ is uniform on t and corresponds to the norm in $\mathcal{L}(L^2)$. Then we get:

$$(78) \quad \|\mathcal{U}_t - e^{-itH_1} \mathcal{U}_0\| = O(h^\infty).$$

□

Lemma 10. *Under the notations used above, we have*

$$(79) \quad l_1^\pm \alpha_j r_1^\pm = l_2^\pm \alpha_j r_2^\pm, \quad l_1^\pm \alpha_j r_2^\pm = l_2^\pm \alpha_j r_1^\pm = 0, \quad j = 1, 2, 3.$$

Proof. As Rubinow and Keller in [37] let us work in a general situation.

We consider the n Hermitian matrices M_μ and n real scalars p_μ , $\mu = 1, \dots, n$. Let G be the Hermitian matrix defined by

$$G = \sum_{\mu=1}^n p_\mu M_\mu.$$

Let λ be a multiple eigenvalue of G and $B_1, \dots, B_q$, a set of corresponding orthonormal eigenvectors which are differentiable functions of p_μ . Then

$$(80) \quad B_j^\dagger B_k = \delta_{jk},$$

$$(81) \quad G B_k = \lambda B_k.$$

The subscript $\dagger$ designates the complex conjugate of the transpose.

If $\lambda(p_1, \dots, p_\mu)$ is differentiable, we differentiate (81) with respect to p_μ and obtain

$$(82) \quad M_\mu B_k + G \frac{\partial B_k}{\partial p_\mu} = \frac{\partial \lambda}{\partial p_\mu} B_k + \lambda \frac{\partial B_k}{\partial p_\mu}.$$

The multiplication of (82) on the left by $B_j^\dagger$, the use of (81), and the fact that G is Hermitian yield

$$(83) \quad B_j^\dagger M_\mu B_k = \frac{\partial \lambda}{\partial p_\mu} \delta_{jk}.$$

In order to treat our case, we take

$$G = \mathcal{M}^\pm = \sum_{\mu=1}^5 p_\mu^\pm M_\mu,$$

where $M_j = \alpha_j$ for $j = 1, 2, 3$, $M_4 = \beta$ and $M_5 = I_4$ are Hermitian matrices (α_j, β are the Dirac matrices) and $p_\mu^\pm$ are five real scalars.

We also take $\lambda^\pm = p_5^\pm \pm \sqrt{(p_1^\pm)^2 + (p_2^\pm)^2 + (p_3^\pm)^2 + (p_4^\pm)^2}$ and $F^\pm$ the point which have the coordinate $p_\mu^\pm$: $p_j^\pm = c\partial_{x_j}\Phi^\pm - eA_j$ for $j = 1, 2, 3$, $p_4^\pm = mc^2 + \frac{e(v_+ - v_-)}{2}$, $p_5^\pm = \partial_t\Phi^\pm + \frac{e(v_+ + v_-)}{2}$.

When $\Phi^\pm$ satisfies (66) and (67), $r_1^\pm, r_2^\pm$ are two orthogonal eigenvectors of $\mathcal{M}^\pm$ corresponding to the eigenvalue $\lambda^\pm = \lambda^\pm(F^\pm) = 0$. Since $|e(v_+ - v_-)| < 2mc^2$ (see (9)), $\lambda^\pm$ is differentiable near the point $F^\pm$. Now, we apply (83) with $B_j^\dagger = l_j^\pm$ and $B_k = r_k^\pm$. After the normalization of $r_k^\pm, l_j^\pm$ we obtain

$$l_j^\pm M_\mu r_k^\pm = \frac{\partial \lambda^\pm(p_1^\pm, \dots, p_5^\pm)}{\partial p_\mu^\pm} \Big|_{F^\pm} (\mp 2p_5^\pm u^\pm) \delta_{jk},$$

and we get the lemma. $\square$

Acknowledgments. The author is grateful to V. Bruneau and J.-F. Bony for many helpful discussions. We also thank the French ANR (Grant no. JC0546063) for the partial financial support.

REFERENCES

- [1] J. Aguilar and J. M. Combes, *A class of analytic perturbations for one-body Schrödinger Hamiltonians*. Comm. Math. Phys. 22 (1971), 269–279.
- [2] V. I. Arnold, *Mathematical Methods of Classical Mechanics*, Springer, New York-Heidelberg-Berlin, 1978 (English translation from Russian).
- [3] M. S. Birman and M. G. Krein, *On the theory of wave operators and scattering operators*. (Russian) Dokl. Akad. Nauk SSSR 144 (1962) 475–478.
- [4] M. S. Birman and D. R. Yafaev, *The spectral shift function. The papers of M. G. Kreĭn and their further development*. St. Petersburg Math. J. 4 (1993) 833–870.
- [5] J.-F. Bony, *Minoration du nombre de résonances engendrées par une trajectoire fermée*, Comm. Partial Differential Equation 27 (2002), 1021–1078.
- [6] J.-F. Bony and J. Sjöstrand, *Traceformula for resonances in small domains*. J. Funct. Anal. 184 (2001), no. 2, 402–418.
- [7] V. Bruneau and V. Petkov, *Representation of the spectral shift function and spectral asymptotics for trapping perturbations*. Comm. Partial Differential Equations 26 no. 11–12 (2001), 2081–2119.
- [8] V. Bruneau and V. Petkov, *Meromorphic continuation of the spectral shift function*, Duke Math. J. 116 (2003) 389–430.
- [9] V. Bruneau and V. Petkov, *Eigenvalues of the reference operator and semiclassical resonances*. J. Funct. Anal. 202 (2003), no. 2, 571–590.
- [10] V. Bruneau and D. Robert, *Asymptotics of the scattering phase for the Dirac operator: High energy, semi-classical and non-relativistic limits*. Ark. Mat., 37 (1999), 1–32.
- [11] N. Burq and M. Zworski, *Resonance expansions in semi-classical propagation*. Comm. Math. Phys. 223 no. 1 (2001), 1–12.
- [12] M. Dimassi and J. Sjöstrand, *Spectral asymptotics in the Semi-classical Limit*, London Math. Soc. Lecture Note Ser. 268, Univ. press, Cambridge, 1990.
- [13] M. Dimassi and V. Petkov, *Spectral Shift function and resonances for non-semi-bounded and Stark Hamiltonians*, J. Math. Pures Appl. 82 (2003) 1303–1342.

- [14] P. A. M. Dirac, *A Theory of Electrons and Protons*, Proc. Roy. Soc. A126 (1930) 360.
- [15] Yu. V. Egorov and M. A. Shubin, *Partial Differential Equation I*, Encyclopaedia of Mathematical Sciences, Vol.30, Springer-Verlag (1992).
- [16] C. Gérard, A. Martinez and D. Robert, *Breit-Wigner formulas for the scattering phase and the total scattering cross-section in the semi-classical limit*. Comm. Math. Phys. 121 no. 2 (1989), 323–336.
- [17] A. Grigis and J. Sjöstrand, *Microlocal analysis for differential operators*. Cambridge University Press, 1994. - (London Mathematical Society Lecture Note series; 196).
- [18] B. Helffer and A. Martinez, *Comparaison entre les diverses notions de résonances*. Helv. Phys. Acta 60 (1987), no. 8, 992–1003.
- [19] B. Helffer and D. Robert, *Calcul fonctionnel par la transformation de Mellin et opérateurs admissibles*. J. Funct. Anal. 53 (1983), no. 3, 246–268.
- [20] B. Helffer and J. Sjöstrand, *Résonances en limite semi-classique*. Mém. Soc. Math. France (N.S.) No. 24-25 (1986), iv+228 pp.
- [21] W. Hunziker, *Distortion analyticity and molecular resonance curves*. Ann. Inst. H. Poincaré Phys. Théor. 45 (1986), no. 4, 339–358.
- [22] V. Ivrii, *Microlocal Analysis and Precise Spectral Asymptotics*- Springer Verlag, 1998. - (Springer Monographs in Mathematics).
- [23] T. Kato, *Perturbation Theory*, Springer-Verlag, New York, 1966.
- [24] M. G. Krein, *On the trace formula in perturbation theory*. (Russian) Mat. Sbornik N.S. 33 (75), (1953). 597–626.
- [25] I. M. Lifshits, *On a problem of the theory of perturbations connected with quantum statistics*. (Russian) Uspehi Matem. Nauk (N.S.) 7, (1952). no. 1(47), 171–180.
- [26] R.B. Melrose, *Weyl asymptotics for the phase in obstacle scattering*. Comm. Partial Differential Equations 13 (1988), no. 11, 1431–1439.
- [27] L. Nedelec, *Resonances for matrix Schrödinger operators*. Duke Math. J. 106 (2001), no. 2, 209–236.
- [28] S. Nakamura, *Spectral shift function for trapping energies in the semiclassical limit*. Comm. Math. Phys. 208 (1999), no. 1, 173–193.
- [29] B. Parisse, *Résonances pour l'opérateur de Dirac*. Helv. Phys. Acta 64 (1991), no. 5, 557–591.
- [30] B. Parisse, *Résonances pour l'opérateur de Dirac. II*. Helv. Phys. Acta 65 (1992), no. 8, 1077–1118.
- [31] V. Petkov and M. Zworski, *Breit-Wigner approximation and the distribution of resonances*. Comm. Math. Phys. 204 (1999), no. 2, 329–351.
- [32] V. Petkov and M. Zworski, *Semi-classical estimates on the scattering determinant*. Ann. Henri Poincaré $\frac{1}{2}$ 2 (2001), no. 4, 675–711.
- [33] V. Petkov, *Weyl asymptotic of the scattering phase for metric perturbations*. Asymptotic Anal. 10 (1995), no. 3, 245–261.
- [34] D. Robert, *Autour de l'approximation semi-classic*, Progr.Math.68, Birkhäuser, Boston, 1987.
- [35] D. Robert, *Semiclassical asymptotics for the spectral shift function*. Differential operators and spectral theory, 187–203, Amer. Math. Soc. Transl. Ser. 2, 189, Amer. Math. Soc., Providence, RI, 1999.
- [36] D. Robert, *Relative time-delay for perturbations of elliptic operators and semiclassical asymptotics*. J. Funct. Anal. 126 (1994), no. 1, 36–82.
- [37] S.I. Rubinow and J.B. Keller, *asymptotic solution of the Dirac equation*, Phys. Rev. 131 (1963), 2789–2796.
- [38] J. Sjöstrand, "A trace formula and review of some estimates for resonances" in *Microlocal Analysis and Spectral theory* (Lucca, Italy, 1996), Nato Adv.Sci. Inst. Ser.C Math. Phys. sci 490, Kluwer, Dordrecht, 1997, 377–437.
- [39] J. Sjöstrand, *Resonances for bottles and trace formulae*, Math. Nachr. 221 (2001), 95–149.
- [40] J. Sjöstrand and M. Zworski, *Complex scaling and the distribution of scattering poles*. J. Amer. Math. Soc. 4 (1991), no. 4, 729–769.
- [41] P. Seba, *The complex scaling method for Dirac resonances*. Lett. Math. Phys. 16 (1988), no. 1, 51–59.
- [42] S. H. Tang and M. Zworski, *Resonance expansions of scattered waves*. Comm. Pure Appl. Math. 53 (2000), no. 10, 1305–1334.
- [43] B. Thaller, *The Dirac equation*. - Springer Verlag, 1992. - (Texts and Monographs in Physics).
- [44] D. R. Yafaev, *Mathematical scattering Theory*, Transl. Math.Monogr. 105, Amer. Math.Soc., Providence, 1992.
- [45] D. R. Yafaev, *A trace formula for the Dirac operator*. Bull. London Math. Soc. 37 (2005), no. 6, 908–918.

- [46] K.Yajima: *The quasi-classical approximation to Dirac equation*, I, J. Fac. Sci. Univ. Tokyo 29 (1982), 161-194.

E-mail address: Abdallah.Khochman@math.u-bordeaux1.fr

MATHÉMATIQUES APPLIQUÉES, UNIVERSITÉ BORDEAUX I, 351, COURS DE LA LIBÉRATION, 33405 TALENCE, FRANCE